





# DOC-1R Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-00384
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	IHC;IF;ELISA
<b>Gene Name</b>	CDK2AP2
<b>Protein Name</b>	Cyclin-dependent kinase 2-associated protein 2
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human CDKA2. AA range:51-100
<b>Specificity</b>	DOC-1R Polyclonal Antibody detects endogenous levels of DOC-1R protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	IHC: 1/100 - 1/300. ELISA: 1/5000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	CDK2AP2; DOC1R; Cyclin-dependent kinase 2-associated protein 2; CDK2-associated protein 2; DOC-1-related protein; DOC-1R
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm . Nucleus . Accumulates in immature oocytes in the nucleus. During the first meiotic division, accumulates in the cytoplasm and localizes in dots in the vicinity of the chromosomes in a region enriched in microtubules. .
<b>Tissue Specificity</b>	Ubiquitous.
<b>Function</b>	similarity:Belongs to the CDK2AP family.,tissue specificity:Ubiquitous.,
<b>Background</b>	cyclin dependent kinase 2 associated protein 2(CDK2AP2) Homo sapiens This gene encodes a protein that interacts with cyclin-dependent kinase 2 associated protein 1. Pseudogenes associated with this gene are located on chromosomes 7 and 9. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, Dec 2012],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**

Immunohistochemistry analysis of paraffin-embedded human brain tissue, using CDKA2 Antibody. The picture on the right is blocked with the synthesized peptide.