



MDMX Polyclonal Antibody

Catalog No	YP-Ab-00447
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	MDM4
Protein Name	Protein Mdm4
Immunogen	The antiserum was produced against synthesized peptide derived from human MDM4. AA range:336-385
Specificity	MDMX Polyclonal Antibody detects endogenous levels of MDMX protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MDM4; MDMX; Protein Mdm4; Double minute 4 protein; Mdm2-like p53-binding protein; Protein Mdmx; p53-binding protein Mdm4
Observed Band	45kD
Cell Pathway	Nucleus.
Tissue Specificity	Expressed in all tissues tested with high levels in thymus.
Function	alternative products:Additional isoforms seem to exist,domain:Region I is sufficient for binding p53 and inhibiting its G1 arrest and apoptosis functions. It also binds p73. Region II contains most of a central acidic region and a putative C4-type zinc finger. The RING finger domain which coordinates two molecules of zinc mediates the heterooligomerization with MDM2.,function:Inhibits p53- and p73-mediated cell cycle arrest and apoptosis by binding its transcriptional activation domain. Inhibits degradation of MDM2. Can reverse MDM2-targeted degradation of p53 while maintaining suppression of p53 transactivation and apoptotic functions.,mass spectrometry: PubMed:11840567,similarity:Belongs to the MDM2/MDM4 family.,similarity:Contains 1 RanBP2-type zinc finger.,similarity:Contains 1 RING-type zinc finger.,similarity:Contains 1 SWIB domain.,subunit:Binds to p53, p73 and MDM2.,tissue specif

**Background**

This gene encodes a nuclear protein that contains a p53 binding domain at the N-terminus and a RING finger domain at the C-terminus, and shows structural similarity to p53-binding protein MDM2. Both proteins bind the p53 tumor suppressor protein and inhibit its activity, and have been shown to be overexpressed in a variety of human cancers. However, unlike MDM2 which degrades p53, this protein inhibits p53 by binding its transcriptional activation domain. This protein also interacts with MDM2 protein via the RING finger domain, and inhibits the latter's degradation. So this protein can reverse MDM2-targeted degradation of p53, while maintaining suppression of p53 transactivation and apoptotic functions. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, Feb 2011],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of 293T cells using MDMX Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using MDM4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from rat muscle cells, using MDM4 Antibody. The lane on the right is blocked with the synthesized peptide.