



# ML-IAP Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-00452
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;IHC
<b>Gene Name</b>	BIRC7
<b>Protein Name</b>	Baculoviral IAP repeat-containing protein 7
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human ML-IAP. AA range:162-211
<b>Specificity</b>	ML-IAP Polyclonal Antibody detects endogenous levels of ML-IAP protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000;IHC-p 1:50-300
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	BIRC7; KIAP; LIVIN; MLIAP; RNF50; Baculoviral IAP repeat-containing protein 7; Kidney inhibitor of apoptosis protein; KIAP; Livin; Melanoma inhibitor of apoptosis protein; ML-IAP; RING finger protein 50
<b>Observed Band</b>	21kD
<b>Cell Pathway</b>	Nucleus . Cytoplasm . Golgi apparatus . Nuclear, and in a filamentous pattern throughout the cytoplasm. Full-length livin is detected exclusively in the cytoplasm, whereas the truncated form (tLivin) is found in the peri-nuclear region with marked localization to the Golgi apparatus; the accumulation of tLivin in the nucleus shows positive correlation with the increase in apoptosis.
<b>Tissue Specificity</b>	Isoform 1 and isoform 2 are expressed at very low levels or not detectable in most adult tissues. Detected in adult heart, placenta, lung, lymph node, spleen and ovary, and in several carcinoma cell lines. Isoform 2 is detected in fetal kidney, heart and spleen, and at lower levels in adult brain, skeletal muscle and peripheral blood leukocytes.
<b>Function</b>	function:Protects against apoptosis induced by TNF or by chemical agents such as adriamycin, etoposide or staurosporine. Suppression of apoptosis is mediated by activation of MAPK8/JNK1, and possibly also of MAPK9/JNK2. This activation depends on TAB1 and NR2C2/TAK1. In vitro, inhibits caspase-3 and proteolytic activation of pro-caspase-9. Isoform 1 blocks staurosporine-induced apoptosis and isoform 2 blocks etoposide-induced apoptosis.,similarity:Belongs to the IAP



family.,similarity:Contains 1 BIR repeat.,similarity:Contains 1 RING-type zinc finger.,subcellular location:Nuclear, and in a filamentous pattern throughout the cytoplasm.,subunit:Binds to caspase-9. Interaction with SMAC via the BIR domain disrupts binding to caspase-9 and apoptotic suppressor activity. Interacts with TAB1. In vitro, interacts with caspase-3 and caspase-7 via its BIR domain.,tissue specificity:Very low level

#### Background

This gene encodes a member of the inhibitor of apoptosis protein (IAP) family, and contains a single copy of a baculovirus IAP repeat (BIR) as well as a RING-type zinc finger domain. The BIR domain is essential for inhibitory activity and interacts with caspases, while the RING finger domain sometimes enhances antiapoptotic activity but does not inhibit apoptosis alone. Elevated levels of the encoded protein may be associated with cancer progression and play a role in chemotherapy sensitivity. Alternative splicing results in multiple transcript variants [provided by RefSeq, Jul 2013],

#### matters needing attention


Avoid repeated freezing and thawing!

#### Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




## Products Images



Western Blot analysis of various cells using ML-IAP Polyclonal Antibody



Western blot analysis of lysate from COLO cells, using ML-IAP antibody.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).