



MYH Polyclonal Antibody

Catalog No	YP-Ab-00456
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	MUTYH
Protein Name	A/G-specific adenine DNA glycosylase
Immunogen	The antiserum was produced against synthesized peptide derived from human MUTYH. AA range:151-200
Specificity	MYH Polyclonal Antibody detects endogenous levels of MYH protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MUTYH; MYH; A/G-specific adenine DNA glycosylase; MutY homolog; hMYH
Observed Band	60kD
Cell Pathway	Nucleus . Mitochondrion .
Tissue Specificity	Kidney,
Function	cofactor: Binds 1 4Fe-4S cluster. The cluster is not important for the catalytic activity, but which is probably involved in the proper positioning of the enzyme along the DNA strand.; disease: Defects in MUTYH are a cause of autosomal recessive colorectal adenomatous polyposis [MIM:608456].; disease: Defects in MUTYH are a cause of gastric cancer [MIM:137215].; function: Involved in oxidative DNA damage repair. Initiates repair of A*oxoG to C*G by removing the inappropriately paired adenine base from the DNA backbone. Possesses both adenine and 2-OH-A DNA glycosylase activities.; similarity: Belongs to the nth/mutY family.; similarity: Contains 1 nudix hydrolase domain.;
Background	This gene encodes a DNA glycosylase involved in oxidative DNA damage repair. The enzyme excises adenine bases from the DNA backbone at sites where adenine is inappropriately paired with guanine, cytosine, or 8-oxo-7,8-dihydroguanine, a major oxidatively damaged DNA lesion. The protein



is localized to the nucleus and mitochondria. Mutations in this gene result in heritable predisposition to colon and stomach cancer. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images




Western Blot analysis of various cells using MYH Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunofluorescence analysis of A549 cells, using MUTYH Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa and HepG2 cells, using MUTYH Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Small intestinal stromal tumor. 1, Tris-EDTA, pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200 (4° overnight). 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).