



c-FLIP Polyclonal Antibody

Catalog No	YP-Ab-00565
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	CFLAR
Protein Name	CASP8 and FADD-like apoptosis regulator
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human CFLAR. AA range:1-50
Specificity	c-FLIP Polyclonal Antibody detects endogenous levels of c-FLIP protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CFLAR; CASH; CASP8AP1; CLARP; MRIT; CASP8 and FADD-like apoptosis regulator; Caspase homolog; CASH; Caspase-eight-related protein; Casper; Caspase-like apoptosis regulatory protein;CLARP; Cellular FLICE-like inhibitory protein; c-FLIP; FADD-like antiapoptotic molecule 1; FLAME-1; Inhibitor of FLICE; I-FLICE; MACH-related inducer of toxicity; MRIT; Usurpin
Observed Band	55kD
Cell Pathway	cytoplasm,cytosol,death-inducing signaling complex,CD95 death-inducing signaling complex,membrane raft,riposome,
Tissue Specificity	Widely expressed. Higher expression in skeletal muscle, pancreas, heart, kidney, placenta, and peripheral blood leukocytes. Also detected in diverse cell lines. Isoform 8 is predominantly expressed in testis and skeletal muscle.
Function	domain:The caspase domain lacks the active sites residues involved in catalysis.,function:Apoptosis regulator protein which may function as a crucial link between cell survival and cell death pathways in mammalian cells. Acts as an inhibitor of TNFRSF6 mediated apoptosis. A proteolytic fragment (p43) is likely retained in the death-inducing signaling complex (DISC) thereby blocking further recruitment and processing of caspase-8 at the complex. Full length and shorter isoforms have been shown either to induce apoptosis or to reduce TNFRSF-triggered apoptosis. Lacks enzymatic (caspase)



activity.,induction:Repressed by IL-2 after TCR stimulation, during progression to the S-phase of the cell cycle.,PTM:Proteolytically processed; probably by caspase-8. Processing likely occurs at the DISC and generates subunit p43 and p12.,similarity:Belongs to the peptidase C14A family.,similarity:Contains

Background

The protein encoded by this gene is a regulator of apoptosis and is structurally similar to caspase-8. However, the encoded protein lacks caspase activity and appears to be itself cleaved into two peptides by caspase-8. Several transcript variants encoding different isoforms have been found for this gene, and partial evidence for several more variants exists. [provided by RefSeq, Feb 2011],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of SW480, SKOV3, HepG2 cells using c-FLIP Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from SW480 cells, using CFLAR Antibody.