



ATP7B Polyclonal Antibody

Catalog No	YP-Ab-00678
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	ATP7B
Protein Name	Copper-transporting ATPase 2
Immunogen	The antiserum was produced against synthesized peptide derived from human ATP7B. AA range:161-210
Specificity	ATP7B Polyclonal Antibody detects endogenous levels of ATP7B protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ATP7B; PWD; WC1; WND; Copper-transporting ATPase 2; Copper pump 2; Wilson disease-associated protein
Observed Band	
Cell Pathway	Golgi apparatus, trans-Golgi network membrane ; Multi-pass membrane protein . Late endosome . Predominantly found in the trans-Golgi network (TGN). Localized in the trans-Golgi network under low copper conditions, redistributes to cytoplasmic vesicles when cells are exposed to elevated copper levels, and then recycles back to the trans-Golgi network when copper is removed (PubMed:10942420). .; [Isoform 1]: Golgi apparatus membrane ; Multi-pass membrane protein .; [Isoform 2]: Cytoplasm .; [WND/140 kDa]: Mitochondrion .
Tissue Specificity	Most abundant in liver and kidney and also found in brain. Isoform 2 is expressed in brain but not in liver. The cleaved form WND/140 kDa is found in liver cell lines and other tissues.
Function	catalytic activity:ATP + H(2)O + Cu(2+)(In) = ADP + phosphate + Cu(2+)(Out).,disease:Defects in ATP7B are the cause of Wilson disease (WD) [MIM:277900]. WD is an autosomal recessive disorder of copper metabolism in which copper cannot be incorporated into ceruloplasmin in liver, and cannot be excreted from the liver into the bile. Copper accumulates in the liver and subsequently in the brain and kidney. The disease is characterized by neurologic manifestations and signs of cirrhosis.,function:Involved in the export of copper out of the cells, such as the efflux of hepatic copper into the bile.,online



information: Wilson's disease website, PTM: Isoform 1 may be proteolytically cleaved at the N-terminus to produce the WND/140 kDa form., similarity: Belongs to the cation transport ATPase (P-type) family., similarity: Belongs to the cation transport ATPase (P-type) family. Type IB subfamily., simila

Background

This gene is a member of the P-type cation transport ATPase family and encodes a protein with several membrane-spanning domains, an ATPase consensus sequence, a hinge domain, a phosphorylation site, and at least 2 putative copper-binding sites. This protein functions as a monomer, exporting copper out of the cells, such as the efflux of hepatic copper into the bile. Alternate transcriptional splice variants, encoding different isoforms with distinct cellular localizations, have been characterized. Mutations in this gene have been associated with Wilson disease (WD). [provided by RefSeq, Jul 2008],



matters needing attention

Avoid repeated freezing and thawing!



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunofluorescence analysis of HeLa cells, using ATP7B Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human testis tissue, using ATP7B Antibody. The picture on the right is blocked with the synthesized peptide.