



# eEF1A2 binding protein Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-01212
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;Pig
<b>Applications</b>	WB;IHC;IF
<b>Gene Name</b>	IGFN1
<b>Protein Name</b>	Immunoglobulin-like and fibronectin type III domain-containing protein 1 (EEF1A2-binding protein 1) (KY-interacting protein 1)
<b>Immunogen</b>	Recombinant Protein of eEF1A2 binding protein
<b>Specificity</b>	The antibody detects endogenous eEF1A2 binding protein protein
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB: 1:500-1:2000 IHC: 1:50-1:200. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Immunoglobulin-like and fibronectin type III domain-containing protein 1 (EEF1A2-binding protein 1;KY-interacting protein 1)
<b>Observed Band</b>	115kD
<b>Cell Pathway</b>	Nucleus . Cytoplasm, myofibril, sarcomere, Z line .
<b>Tissue Specificity</b>	Expressed in skeletal muscle.
<b>Function</b>	similarity:Contains 4 fibronectin type-III domains.,similarity:Contains 5 Ig-like (immunoglobulin-like) domains.,subunit:Interacts with KY.,tissue specificity:Expressed in skeletal muscle.,
<b>Background</b>	similarity:Contains 4 fibronectin type-III domains.,similarity:Contains 5 Ig-like (immunoglobulin-like) domains.,subunit:Interacts with KY.,tissue specificity:Expressed in skeletal muscle.,
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images

Western blot analysis of Pig Skeletal Muscle with eEF1A2 Rabbit pAb diluted at 1:1,000

130KD —  
95KD —  
70KD —  
55KD —



Immunohistochemical analysis of paraffin-embedded Pig Skeletal Muscle using eEF1A2Rabbit pAb diluted at 1:100

