



DDX54 Polyclonal Antibody

Catalog No	YP-Ab-01646
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	DDX54
Protein Name	ATP-dependent RNA helicase DDX54
Immunogen	Synthesized peptide derived from DDX54 . at AA range: 570-650
Specificity	DDX54 Polyclonal Antibody detects endogenous levels of DDX54 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DDX54; ATP-dependent RNA helicase DDX54; ATP-dependent RNA helicase DP97; DEAD box RNA helicase 97 kDa; DEAD box protein 54
Observed Band	
Cell Pathway	Nucleus, nucleolus .
Tissue Specificity	Expressed in the fallopian tube, cervix and uterus. Also expressed in the brain.
Function	caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data.,function:Has RNA-dependent ATPase activity. Represses the transcriptional activity of nuclear receptors.,similarity:Belongs to the DEAD box helicase family. DDX54/DBP10 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,subunit:Interacts in a hormone-dependent manner with nuclear receptors.,
Background	This gene encodes a member of the DEAD box protein family. DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division.



The nucleolar protein encoded by this gene interacts in a hormone-dependent manner with nuclear receptors, and represses their transcriptional activity. Alternative splice variants that encode different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of various cells using DDX54 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).