

(Tel: 400-999-8863 ■ Email:Upingbio.163.com





Helios Polyclonal Antibody

IKZF2. AA range:331-380 Specificity Helios Polyclonal Antibody detects endogenous levels of Helios protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2 Observed Band 56kD Cell Pathway Nucleus. Tissue Specificity Epithelium, Testis, Function function: Associates with Ikaros at centromeric heterochromatin., similarity: Beld to the Ikaros C2H2-type zinc-finger protein family, similarity: Contains 6 C2H2-zinc fingers., subunit: Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. Thromembers of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with othe Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding of some variants encoding of some variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding of some variants encodin		
Applications WB;ELISA Gene Name IKZF2 Protein Name Immunogen The antiserum was produced against synthesized peptide derived from huma IKZF2. AA range:331-380 Specificity Helios Polyclonal Antibody detects endogenous levels of Helios protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finge protein 2 Observed Band Cell Pathway Nucleus . Tissue Specificity Epithelium, Testis, Function function: Associates with Ikaros at centromeric heterochromatin, similarity. Belc to the Ikaros C2H2-type zinc-finger protein family, similarity. Contains 6 C2H2-zinc fingers, subunit: Interacts with IKZF4 AND IKZF5. This gene encodes a member of the Ikaros family of zinc-finger proteins. Thr members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with oth Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some varient in the protein family and part of the spene, but the biological validity of some varient in the protein family and part variants encoding different isoforms have been found for this gene, but the biological validity of some varient in the protein family and part variants encoding different in soforms have been found for this gene, but the biological validity of some varients.	Catalog No	YP-Ab-01770
Applications WB;ELISA Gene Name IKZF2 Protein Name Zinc finger protein Helios Immunogen The antiserum was produced against synthesized peptide derived from huma IKZF2. AA range:331-380 Specificity Helios Polyclonal Antibody detects endogenous levels of Helios protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2 Observed Band 56kD Cell Pathway Nucleus. Tissue Specificity Epithelium,Testis, function: Associates with Ikaros at centromeric heterochromatin, similarity:Beld to the Ikaros C2H2-type zinc-finger protein family. similarity:Contains 6 C2H2-zinc fingers, subunit:Interacts with IKZF4 AND IKZF5. Background This gene encodes a member of the Ikaros family of zinc-finger proteins. Thr members of this protein family (Ikaros, Aiolos and Helios) are hematopoletic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with oth Ikaros family members, and is thought to function predominaty in early hematopoletic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Isotype	IgG
Gene Name IKZF2 Protein Name Zinc finger protein Helios Immunogen The antiserum was produced against synthesized peptide derived from huma IKZF2. AA range:331-380 Specificity Helios Polyclonal Antibody detects endogenous levels of Helios protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finge protein 2 Observed Band 56kD Cell Pathway Nucleus . Function Epithelium,Testis, Function This gene encodes a member of the Ikaros family of zinc-finger proteins family, similarity. Contains 6 C2H2-zinc fingers, subunit Interacts with IKZF4 AND IKZF5. Background This gene encodes a member of the Ikaros family of zinc-finger proteins. Thremembers of this protein family (Ikaros, Alolos and Helios) are hematopoletic-speci	Reactivity	Human;Mouse;Rat
Protein Name Zinc finger protein Helios Immunogen The antiserum was produced against synthesized peptide derived from human IKZF2. AA range:331-380 Specificity Helios Polyclonal Antibody detects endogenous levels of Helios protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finge protein 2 Observed Band 56kD Cell Pathway Nucleus . Function function:Associates with Ikaros at centromeric heterochromatin., similarity:Beld to the Ikaros C2H2-type zinc-finger protein family, willingers., subunit: interacts with IKZF4 AND IKZF5. Background This gene encodes a member of the Ikaros family of zinc-finger proteins. Thremembers of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms home- or hetero-dimers w	Applications	WB;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from human IKZF2. AA range:331-380 Specificity Helios Polyclonal Antibody detects endogenous levels of Helios protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finge protein 2 Observed Band 56kD Cell Pathway Nucleus . Function Epithelium, Testis, Function function:Associates with Ikaros at centromeric heterochromatin., similarity:Beld to the Ikaros C2H2-type zinc-finger protein family, ismilarity:Contains 6 C2H2-zinc fingers, subunit:Interacts with IKZF4 AND IKZF5. Background This gene encodes a member of the Ikaros family of zinc-finger proteins. Thremembers of this protein family, (karos, Alolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms home-or	Gene Name	IKZF2
IKZF2. AA range:331-380 Specificity Helios Polyclonal Antibody detects endogenous levels of Helios protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2 Observed Band 56kD Cell Pathway Nucleus. Tissue Specificity Epithelium,Testis, Function function:Associates with Ikaros at centromeric heterochromatin.,similarity:Beld to the Ikaros C2H2-type zinc-finger protein family,similarity:Contains 6 C2H2-zinc fingers.,subunit:Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. Thromebers of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with othe Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding of some	Protein Name	Zinc finger protein Helios
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finge protein 2 Observed Band Cell Pathway Nucleus. Tissue Specificity Epithelium, Testis, Function function: Associates with Ikaros at centromeric heterochromatin., similarity: Beld to the Ikaros C2H2-type zinc-finger protein family, similarity: Contains 6 C2H2-zinc fingers, subunit:Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. The members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with othe Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different is forms have been found for this gene, but the biological validity of some vari	Immunogen	The antiserum was produced against synthesized peptide derived from human IKZF2. AA range:331-380
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2 Observed Band 56kD Cell Pathway Nucleus . Tissue Specificity Epithelium,Testis, Function function:Associates with Ikaros at centromeric heterochromatin., similarity:Beld to the Ikaros C2H2-type zinc-finger protein family., similarity:Contains 6 C2H2-zinc fingers, subunit:Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. Thr members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with othe Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some variant senced in gene, but the biological validity of some variants and control in the regulation of processor in the protein forms homo- or hetero-dimers with othe sorem hav	Specificity	Helios Polyclonal Antibody detects endogenous levels of Helios protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2 Observed Band 56kD Cell Pathway Nucleus. Tissue Specificity Epithelium,Testis, Function function:Associates with Ikaros at centromeric heterochromatin.,similarity:Beld to the Ikaros C2H2-type zinc-finger protein family, similarity:Contains 6 C2H2-zinc fingers.,subunit:Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with othe Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding different isoforms have been found for this gene, but the biological validity of some variants encoding different isoforms have been found for this gene, but the biological validity of some variants.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finge protein 2 Observed Band 56kD Cell Pathway Nucleus . Tissue Specificity Epithelium, Testis, Function function: Associates with Ikaros at centromeric heterochromatin., similarity: Beld to the Ikaros C2H2-type zinc-finger protein family, similarity: Contains 6 C2H2-zinc fingers., subunit: interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. The members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with othe Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Source	Polyclonal, Rabbit,IgG
applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2 Observed Band 56kD Cell Pathway Nucleus. Tissue Specificity Epithelium, Testis, Function function: Associates with Ikaros at centromeric heterochromatin., similarity: Belot to the Ikaros C2H2-type zinc-finger protein family., similarity: Contains 6 C2H2-zinc fingers., subunit: Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. The members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with othe Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2 Observed Band 56kD Cell Pathway Nucleus . Tissue Specificity Epithelium, Testis, Function function: Associates with Ikaros at centromeric heterochromatin., similarity: Belot to the Ikaros C2H2-type zinc-finger protein family, similarity: Contains 6 C2H2-zinc fingers., subunit: Interacts with IkZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. The members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other likaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Dilution	
Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2 Observed Band 56kD Cell Pathway Nucleus . Tissue Specificity Epithelium, Testis, Function function: Associates with Ikaros at centromeric heterochromatin., similarity: Below to the Ikaros C2H2-type zinc-finger protein family., similarity: Contains 6 C2H2-zinc fingers., subunit: Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. The members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other laterals and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Concentration	1 mg/ml
Synonyms IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2 Observed Band 56kD Cell Pathway Nucleus . Epithelium, Testis, Function function: Associates with Ikaros at centromeric heterochromatin., similarity: Belot to the Ikaros C2H2-type zinc-finger protein family., similarity: Contains 6 C2H2-zinc fingers., subunit: Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. The members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other larger protein family (Ikaros, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Purity	≥90%
Doserved Band Cell Pathway Nucleus . Tissue Specificity Epithelium, Testis, function function: Associates with Ikaros at centromeric heterochromatin., similarity: Belot to the Ikaros C2H2-type zinc-finger protein family., similarity: Contains 6 C2H2-zinc fingers., subunit: Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. The members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other largest protein family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Storage Stability	-20°C/1 year
Tissue Specificity Epithelium, Testis, function function: Associates with Ikaros at centromeric heterochromatin., similarity: Belove to the Ikaros C2H2-type zinc-finger protein family., similarity: Contains 6 C2H2-zinc fingers., subunit: Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. The members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other likaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Synonyms	IKZF2; HELIOS; ZNFN1A2; Zinc finger protein Helios; Ikaros family zinc finger protein 2
Tissue Specificity Epithelium, Testis, function function: Associates with Ikaros at centromeric heterochromatin., similarity: Belove to the Ikaros C2H2-type zinc-finger protein family., similarity: Contains 6 C2H2-zinc fingers., subunit: Interacts with IKZF4 AND IKZF5., Background This gene encodes a member of the Ikaros family of zinc-finger proteins. The members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other likaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some variance.	Observed Band	56kD
function: Associates with Ikaros at centromeric heterochromatin., similarity: Beld to the Ikaros C2H2-type zinc-finger protein family., similarity: Contains 6 C2H2-zinc fingers., subunit: Interacts with IKZF4 AND IKZF5., This gene encodes a member of the Ikaros family of zinc-finger proteins. Thr members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other likaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some variance.	Cell Pathway	Nucleus .
to the Ikaros C2H2-type zinc-finger protein family.,similarity:Contains 6 C2H2-zinc fingers.,subunit:Interacts with IKZF4 AND IKZF5., This gene encodes a member of the Ikaros family of zinc-finger proteins. Thr members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other likaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Tissue Specificity	Epithelium,Testis,
members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with othe Ikaros family members, and is thought to function predominantly in early hematopoietic development. Multiple transcript variants encoding different isoforms have been found for this gene, but the biological validity of some vari	Function	function: Associates with Ikaros at centromeric heterochromatin., similarity: Belongs to the Ikaros C2H2-type zinc-finger protein family., similarity: Contains 6 C2H2-type zinc fingers., subunit: Interacts with IKZF4 AND IKZF5.,
has not been determined. [provided by ReiSeq, Jul 2000],	Background	This gene encodes a member of the Ikaros family of zinc-finger proteins. Three members of this protein family (Ikaros, Aiolos and Helios) are hematopoietic-specific transcription factors involved in the regulation of lymphocyte development. This protein forms homo- or hetero-dimers with other Ikaros family members, and is thought to function predominantly in early



UpingBio technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of Hela cells using Helios Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Western blot analysis of the lysates from HeLa cells using IKZF2 antibody.