



Hexim1 Polyclonal Antibody

Catalog No	YP-Ab-01774
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	HEXIM1
Protein Name	Protein HEXIM1
Immunogen	The antiserum was produced against synthesized peptide derived from human HEXIM1. AA range:181-230
Specificity	Hexim1 Polyclonal Antibody detects endogenous levels of Hexim1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HEXIM1; CLP1; EDG1; HIS1; MAQ1; Protein HEXIM1; Cardiac lineage protein 1; Estrogen down-regulated gene 1 protein; Hexamethylene bis-acetamide-inducible protein 1; Menage a quatre protein 1
Observed Band	40kD
Cell Pathway	Nucleus . Cytoplasm . Binds alpha-importin and is mostly nuclear (PubMed:16362050).
Tissue Specificity	Ubiquitously expressed with higher expression in placenta. HEXIM1 and HEXIM2 are differentially expressed. Expressed in endocrine tissues.
Function	domain:The coiled-coil domain mediates oligomerization.,function:Transcriptional regulator which functions as a general RNA polymerase II transcription inhibitor. In cooperation with 7SK snRNA sequesters P-TEFb in a large inactive 7SK snRNP complex preventing RNA polymerase II phosphorylation and subsequent transcriptional elongation. May also regulate NF-kappa-B, ESR1, NR3C1 and C/EBP-dependent transcriptional activity.,induction:Up-regulated by HMBA (hexamethylene bisacetamide) (at protein level). Down-regulated by estrogen.,miscellaneous:Inhibits Tat activity which is required for HIV-1 transcription.,similarity:Belongs to the HEXIM family.,subcellular location:Binds alpha-importin and is mostly nuclear.,subunit:Homooligomer and heterooligomer with HEXIM2; probably dimeric. Component of the 7SK snRNP complex at least



composed of P-TEFb (composed of CDK9 and CCNT1/cyclin-T1), HEXIM1, H

Background

Expression of this gene is induced by hexamethylene-bis-acetamide in vascular smooth muscle cells. This gene has no introns. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images




Western Blot analysis of RAT-HEART cells using Hexim1 Polyclonal Antibody diluted at 1:2000




Immunofluorescence analysis of HeLa cells, using HEXIM1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from NIH/3T3 cells, using HEXIM1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HeLa cells using HEXIM1 antibody.



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).