



HIF-1α Polyclonal Antibody

Catalog No	YP-Ab-01776
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IF;WB;IHC;IP;ELISA
Gene Name	HIF1A
Protein Name	Hypoxia-inducible factor 1-alpha
Immunogen	The antiserum was produced against synthesized peptide derived from human HIF-1alpha. AA range:328-377
Specificity	HIF-1α Polyclonal Antibody detects endogenous levels of HIF-1α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IF: 1:50-200 Western Blot: 1/500 - 1/2000.IP 1:200 Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	HIF1A; BHLHE78; MOP1; PASD8; Hypoxia-inducible factor 1-alpha; HIF-1-alpha; HIF1-alpha; ARNT-interacting protein; Basic-helix-loop-helix-PAS protein MOP1; Class E basic helix-loop-helix protein 78; bHLHe78; Member of PAS protein 1; PAS doma
Observed Band	92-130kD
Cell Pathway	Cytoplasm . Nucleus . Nucleus speckle . Colocalizes with HIF3A in the nucleus and speckles (By similarity). Cytoplasmic in normoxia, nuclear translocation in response to hypoxia (PubMed:9822602). .
Tissue Specificity	Expressed in most tissues with highest levels in kidney and heart. Overexpressed in the majority of common human cancers and their metastases, due to the presence of intratumoral hypoxia and as a result of mutations in genes encoding oncoproteins and tumor suppressors. A higher level expression seen in pituitary tumors as compared to the pituitary gland.
Function	domain:Contains two independent C-terminal transactivation domains, NTAD and CTAD, which function synergistically. Their transcriptional activity is repressed by an intervening inhibitory domain (ID).,function:Functions as a master transcriptional regulator of the adaptive response to hypoxia. Under hypoxic conditions activates the transcription of over 40 genes, including, erythropoietin, glucose transporters, glycolytic enzymes, vascular endothelial growth factor, and other genes whose protein products increase oxygen delivery or facilitate



metabolic adaptation to hypoxia. Plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Binds to core DNA sequence 5'-[AG]CGTG-3' within the hypoxia response element (HRE) of target gene promoters. Activation requires recruitment of transcriptional coactivators such as CREBPB and EP300. Acti

Background

hypoxia inducible factor 1 alpha subunit(HIF1A) Homo sapiens This gene encodes the alpha subunit of transcription factor hypoxia-inducible factor-1 (HIF-1), which is a heterodimer composed of an alpha and a beta subunit. HIF-1 functions as a master regulator of cellular and systemic homeostatic response to hypoxia by activating transcription of many genes, including those involved in energy metabolism, angiogenesis, apoptosis, and other genes whose protein products increase oxygen delivery or facilitate metabolic adaptation to hypoxia. HIF-1 thus plays an essential role in embryonic vascularization, tumor angiogenesis and pathophysiology of ischemic disease. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jul 2011],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images




Western blot analysis of lysates from 1)Hela cell ,
2)Hela cells knockdown by siRNA
(F:GCCACAUUCACGUAUAUGATT,R:UCAUAUACG
UGAAUGUGGCTT), (Green) primary antibody was
diluted at 1:1000, 4° over night, Dylight 800 secondary
antibody(Immunoway:RS23920)was diluted at
1:10000, 37° 1hour. (Red) GAPDH Monoclonal
Antibody(5B7) (Immunoway:YM3029) antibody was
diluted at 1:5000 as loading control, 4° over night,
Dylight 680 secondary
antibody(Immunoway:RS23710)was diluted at
1:10000, 37° 1hour.




Zhao, Shaorong et al. "Deciphering the performance of
polo-like kinase 1 in triple-negative breast cancer
progression according to the centromere protein
U-phosphorylation pathway." American journal of
cancer research vol. 11,5 2142-2158. 15 May. 2021



Zhang, Y., Wang, X., Chu, C. et al. Genetically
engineered magnetic nanocages for cancer
magneto-catalytic theranostics. Nat Commun 11, 5421
(2020).



Jiang, Ruibin, et al. "Total Flavonoids from Carya
cathayensis Sarg. Leaves Alleviate H9c2 Cells
Hypoxia/Reoxygenation Injury via Effects on miR-21
Expression, PTEN/Akt, and the Bcl-2/Bax
Pathway." Evidence-Based Complementary and
Alternative Medicine 2018 (2018).



Han, Dan, et al. "Nickel-smelting fumes increased the expression of HIF-1α through PI3K/ERK pathway in NIH/3T3 cells." *Journal of occupational health* 58.5 (2016): 413-424.