

(Tel: 400-999-8863 ■ Email:Upingbio.163.com





TEF-5 Polyclonal Antibody

| Catalog No | YP-Ab-02097 |
|---------------------------|--|
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB;ELISA |
| Gene Name | TEAD3 |
| Protein Name | Transcriptional enhancer factor TEF-5 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human TEAD3. AA range:211-260 |
| Specificity | TEF-5 Polyclonal Antibody detects endogenous levels of TEF-5 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | TEAD3; TEAD5; TEF5; Transcriptional enhancer factor TEF-5; DTEF-1; TEA domain family member 3; TEAD-3 |
| Observed Band | 49kD |
| Cell Pathway | Nucleus. |
| Tissue Specificity | Preferentially expressed in the placenta. |
| Function | function:Binds to multiple functional elements of the human chorionic somatomammotropin-B gene enhancer.,similarity:Contains 1 TEA DNA-binding domain.,tissue specificity:Preferentially expressed in the placenta., |
| Background | This gene product is a member of the transcriptional enhancer factor (TEF) family of transcription factors, which contain the TEA/ATTS DNA-binding domain. It is predominantly expressed in the placenta and is involved in the transactivation of the chorionic somatomammotropin-B gene enhancer. Translation of this protein is initiated at a non-AUG (AUA) start codon. [provided by RefSeq, Jul 2008], |
| matters needing attention | Avoid repeated freezing and thawing! |
| | |



UpingBio technology Co.,Ltd


(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of the lysates from HUVECcells using TEAD3 antibody.