

(Tel: 400-999-8863 ■ Email:Upingbio.163.com






ZC3H11A Polyclonal Antibody


| Catalog No | YP-Ab-02164 |
|---------------------------|--|
| Isotype | IgG |
| Reactivity | Human;Monkey |
| Applications | WB;ELISA |
| Gene Name | ZC3H11A |
| Protein Name | Zinc finger CCCH domain-containing protein 11A |
| Immunogen | The antiserum was produced against synthesized peptide derived from human ZC3H11A. AA range:291-340 |
| Specificity | ZC3H11A Polyclonal Antibody detects endogenous levels of ZC3H11A protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | ZC3H11A; KIAA0663; ZC3HDC11A; Zinc finger CCCH domain-containing protein 11A |
| Observed Band | 110kD |
| Cell Pathway | Nucleus . |
| Tissue Specificity | Brain, Epithelium, Fetal kidney, Kidney, Placenta, Retina, |
| Function | PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., sequence caution:Contaminating sequence. Potential poly-A sequence., similarity:Contains 3 C3H1-type zinc fingers., |
| Background | PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Contains 3 C3H1-type zinc fingers., |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| | |



Products Images



Western Blot analysis of various cells using ZC3H11A Polyclonal Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysates from COS cells, using ZC3H11A Antibody. The lane on the right is blocked with the synthesized peptide.