




ZC3H13 Polyclonal Antibody


Catalog No	YP-Ab-02165
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	ZC3H13
Protein Name	Zinc finger CCCH domain-containing protein 13
Immunogen	The antiserum was produced against synthesized peptide derived from human ZC3H13. AA range:1571-1620
Specificity	ZC3H13 Polyclonal Antibody detects endogenous levels of ZC3H13 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ZC3H13; KIAA0853; Zinc finger CCCH domain-containing protein 13
Observed Band	200kD
Cell Pathway	Nucleus speckle . Nucleus, nucleoplasm .
Tissue Specificity	Brain,Breast,Epithelium,Fetal brain,T-cell,Testis,Uterus,
Function	PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,
Background	PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Contains 1 C3H1-type zinc finger.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of various cells using ZC3H13 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysates from A549, Jurkat, HepG2 cells, using ZC3H13 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HUVECcells using ZC3H13 antibody.