


(Tel: 400-999-8863 ■ Email:Upingbio.163.com




ABHD8 Polyclonal Antibody

Catalog No	YP-Ab-02459
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;IHC;IF;ELISA
Gene Name	ABHD8
Protein Name	Abhydrolase domain-containing protein 8
Immunogen	The antiserum was produced against synthesized peptide derived from human ABHD8. AA range:291-340
Specificity	ABHD8 Polyclonal Antibody detects endogenous levels of ABHD8 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ABHD8; Abhydrolase domain-containing protein 8
Observed Band	45kD
Cell Pathway	extracellular exosome,
Tissue Specificity	Brain,Embryo,Lymph,Muscle,
Function	similarity:Belongs to the AB hydrolase superfamily.,
	enmany, belongs to the 7.2 Hydronaco capenaniny,
Background	This gene is upstream of, and in a head-to-head orientation with the gene for the mitochondrial ribosomal protein L34. The predicted protein contains alpha/beta hydrolase fold and secretory lipase domains. [provided by RefSeq, Jul 2008],
Background matters needing attention	This gene is upstream of, and in a head-to-head orientation with the gene for the mitochondrial ribosomal protein L34. The predicted protein contains alpha/beta
matters needing	This gene is upstream of, and in a head-to-head orientation with the gene for the mitochondrial ribosomal protein L34. The predicted protein contains alpha/beta hydrolase fold and secretory lipase domains. [provided by RefSeq, Jul 2008],

Products Images



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.



Western blot analysis of lysates from HT-29 and LOVO cells, using ABHD8 Antibody. The lane on the right is blocked with the synthesized peptide.