



COX15 Polyclonal Antibody

Catalog No	YP-Ab-02540
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	COX15
Protein Name	Cytochrome c oxidase assembly protein COX15 homolog
Immunogen	The antiserum was produced against synthesized peptide derived from human COX15. AA range:181-230
Specificity	COX15 Polyclonal Antibody detects endogenous levels of COX15 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/5000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	COX15; Cytochrome c oxidase assembly protein COX15 homolog
Observed Band	
Cell Pathway	Mitochondrion membrane ; Multi-pass membrane protein .
Tissue Specificity	Predominantly found in tissues characterized by high rates of oxidative phosphorylation (OxPhos), including muscle, heart, and brain.
Function	disease:Defects in COX15 are a cause of cytochrome c oxidase deficiency (COX deficiency) [MIM:220110]. COX deficiency is a clinically heterogeneous disorder. The clinical features range from isolated myopathy to severe multisystem disease with onset from infancy to adulthood.,disease:Defects in COX15 are a cause of Leigh syndrome [MIM:256000]. Leigh syndrome is an early-onset progressive neurodegenerative disorder characterized by delayed onset of symptoms, hypotonia, feeding difficulties, failure to thrive, motor regression and brainstem signs. Diagnosis is confirmed by the presence of focal, bilateral lesions in one or more areas of the central nervous system including the brainstem, thalamus, basal ganglia, cerebellum and spinal cord.,function:May be involved in the biosynthesis of heme A.,pathway:Porphyryn metabolism; heme A biosynthesis; heme A from heme O: step 1/1.,similarity:Belo
Background	Cytochrome c oxidase (COX), the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to



oxygen. This component is a heteromeric complex consisting of 3 catalytic subunits encoded by mitochondrial genes and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, and the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes a protein which is not a structural subunit, but may be essential for the biogenesis of COX formation and may function in the hydroxylation of heme O, according to the yeast mutant studies. This protein is predicted to contain 5 transmembrane domains localized in the mitochondrial inner membrane. Alternative splicing of this gene generates two transcript variants diverging


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using COX15 Antibody. The picture on the right is blocked with the synthesized peptide.