



CROT Polyclonal Antibody

Catalog No	YP-Ab-02554
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	CROT
Protein Name	Peroxisomal carnitine O-octanoyltransferase
Immunogen	Synthesized peptide derived from CROT . at AA range: 190-270
Specificity	CROT Polyclonal Antibody detects endogenous levels of CROT protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CROT; COT; Peroxisomal carnitine O-octanoyltransferase; COT
Observed Band	72kD
Cell Pathway	Peroxisome .
Tissue Specificity	Brain,Skin,
Function	catalytic activity:Octanoyl-CoA + L-carnitine = CoA + L-octanoylcarnitine..function:Beta-oxidation of fatty acids. The highest activity concerns the C6 to C10 chain length substrate. Converts the end product of pristanic acid beta oxidation, 4,8-dimethylnonanoyl-CoA, to its corresponding carnitine ester..pathway:Lipid metabolism; fatty acid beta-oxidation..similarity:Belongs to the carnitine/choline acetyltransferase family..subunit:Monomer.,
Background	This gene encodes a member of the carnitine/choline acetyltransferase family. The encoded protein converts 4,8-dimethylnonanoyl-CoA to its corresponding carnitine ester. This transesterification occurs in the peroxisome and is necessary for transport of medium- and long- chain acyl-CoA molecules out of the peroxisome to the cytosol and mitochondria. The protein thus plays a role in lipid metabolism and fatty acid beta-oxidation. Alternatively spliced transcript variants have been described.[provided by RefSeq, Jan 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).