

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com






TPH1 Polyclonal Antibody

Catalog No	YP-Ab-02801
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	TPH1
Protein Name	Tryptophan 5-hydroxylase 1
Immunogen	The antiserum was produced against synthesized peptide derived from human Tryptophan Hydroxylase. AA range:26-75
Specificity	TPH1 Polyclonal Antibody detects endogenous levels of TPH1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TPH1; TPH; TPRH; TRPH; Tryptophan 5-hydroxylase 1; Tryptophan 5-monooxygenase 1
Observed Band	51kD
Cell Pathway	cytosol,neuron projection,
Tissue Specificity	
	[Isoform 2]: Seems to be less widely expressed than isoform 1.
Function	[Isoform 2]: Seems to be less widely expressed than isoform 1. catalytic activity:L-tryptophan + tetrahydrobiopterin + O(2) = 5-hydroxy-L-tryptophan + 4a-hydroxytetrahydrobiopterin.,cofactor:Fe(2+) ion.,pathway:Aromatic compound metabolism; serotonin biosynthesis; serotonin from L-tryptophan: step 1/2.,similarity:Belongs to the biopterin-dependent aromatic amino acid hydroxylase family.,similarity:Contains 1 ACT domain.,subunit:Multimer of identical subunits.,tissue specificity:Isoform 2 seems to be less widely expressed than isoform 1.,
	catalytic activity:L-tryptophan + tetrahydrobiopterin + O(2) = 5-hydroxy-L-tryptophan + 4a-hydroxytetrahydrobiopterin.,cofactor:Fe(2+) ion.,pathway:Aromatic compound metabolism; serotonin biosynthesis; serotonin from L-tryptophan: step 1/2.,similarity:Belongs to the biopterin-dependent aromatic amino acid hydroxylase family.,similarity:Contains 1 ACT domain.,subunit:Multimer of identical subunits.,tissue specificity:Isoform 2 seems



UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 💌 Email:Upingbio.163.com



disorder, suicidal behavior, addictions, and others.[provided by RefSeq, Apr 2009],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Products Images




Western Blot analysis of various cells using TPH1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunofluorescence analysis of HepG2 cells, using Tryptophan Hydroxylase Antibody. The picture on the right is blocked with the synthesized peptide.






Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Tryptophan Hydroxylase Antibody. The picture on the right is blocked with the synthesized peptide.



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com





Western blot analysis of lysates from HepG2 cells, using Tryptophan Hydroxylase Antibody. The lane on the right is blocked with the synthesized peptide.