

Tel: 400-999-8863 
■ Email:Upingbio.163.com





## Amylase Polyclonal Antibody

Catalog No	YP-Ab-02858
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF;ELISA
Gene Name	AMY1A/AMY1B/AMY1C/AMY2A/AMY2B
Protein Name	Alpha-amylase 1/Pancreatic alpha-amylase/Alpha-amylase 2B
Immunogen	The antiserum was produced against synthesized peptide derived from the N-terminal region of human AMY1/2. AA range:61-110
Specificity	Amylase Polyclonal Antibody detects endogenous levels of Amylase protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC-p: 1:100-300 ELISA: 1/20000 IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AMY1A; AMY1; AMY1B; AMY1; AMY1C; AMY1; Alpha-amylase 1; 1,4-alpha-D-glucan glucanohydrolase 1; Salivary alpha-amylase; AMY2A; Pancreatic alpha-amylase; PA;1,4-alpha-D-glucan glucanohydrolase; AMY2B; Alpha-amylase 2B; 1,4-alpha-D-glucan glucanohydrolase 2B; Carcinoid alpha-amylase
Observed Band	60kD
Cell Pathway	extracellular space,extracellular exosome,
Tissue Specificity	Glandular pool- thyroid,Saliva,Thyroid,
Function	catalytic activity:Endohydrolysis of (1->4)-alpha-D-glucosidic linkages in oligosaccharides and polysaccharides.,cofactor:Binds 1 calcium ion per subunit.,cofactor:Binds 1 chloride ion per subunit.,online information:Amylase entry,similarity:Belongs to the glycosyl hydrolase 13 family.,subunit:Monomer.,
Background	Amylases are secreted proteins that hydrolyze 1,4-alpha-glucoside bonds in oligosaccharides and polysaccharides, and thus catalyze the first step in digestior of dietary starch and glycogen. The human genome has a cluster of several amylase genes that are expressed at high levels in either salivary gland or pancreas. This gene encodes an amylase isoenzyme produced by the salivary



## UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 🗷 Emall:Upingbio.163.com



gland. Alternative splicing results in multiple transcript variants encoding the same protein. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


## **Products Images**



Western Blot analysis of HeLa cells using Amylase Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded rat-liver, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded rat-liver, antibody was diluted at 1:100




## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com





Immunohistochemical analysis of paraffin-embedded mouse-liver, antibody was diluted at 1:100



Western blot analysis of lysate from HeLa cells, using AMY1/2 Antibody.