



Myosin Id Polyclonal Antibody

Catalog No	YP-Ab-03164
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;ELISA
Gene Name	MYO1D
Protein Name	Unconventional myosin-Id
Immunogen	The antiserum was produced against synthesized peptide derived from human MYO1D. AA range:825-874
Specificity	Myosin Id Polyclonal Antibody detects endogenous levels of Myosin Id protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MYO1D; KIAA0727; Unconventional myosin-Id
Observed Band	
Cell Pathway	Cytoplasm . Perikaryon . Cell projection, dendrite . Early endosome . Cytoplasm, cell cortex . Colocalizes with the actin cytoskeleton in the cell cortex close to the apical cell membrane. Colocalizes with cytoplasmic puncta that are reminiscent of transport vesicles. .
Tissue Specificity	Expressed in many tissues. Highest levels in brain, followed by lung and ovary; expression is lowest in spleen.
Function	function:Myosins are actin-based motor molecules with ATPase activity. Unconventional myosins serve in intracellular movements. Their highly divergent tails are presumed to bind to membranous compartments, which would be moved relative to actin filaments.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 2 IQ domains.,subunit:Binds calmodulin through its IQ motifs.,tissue specificity:Expressed in many tissues. Highest levels in brain, followed by lung and ovary; expression is lowest in spleen.,
Background	function:Myosins are actin-based motor molecules with ATPase activity. Unconventional myosins serve in intracellular movements. Their highly divergent tails are presumed to bind to membranous compartments, which would be moved relative to actin filaments.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 2 IQ domains.,subunit:Binds calmodulin through its IQ

motifs.,tissue specificity:Expressed in many tissues. Highest levels in brain, followed by lung and ovary; expression is lowest in spleen.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using MYO1D Antibody. The picture on the right is blocked with the synthesized peptide.