



# APS Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-03707
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	SH2B2
<b>Protein Name</b>	SH2B adapter protein 2
<b>Immunogen</b>	Synthesized peptide derived from the Internal region of human APS.
<b>Specificity</b>	APS Polyclonal Antibody detects endogenous levels of APS protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	SH2B2; APS; SH2B adapter protein 2; Adapter protein with pleckstrin homology and Src homology 2 domains; SH2 and PH domain-containing adapter protein APS
<b>Observed Band</b>	67kD
<b>Cell Pathway</b>	Cytoplasm . Cell membrane . Cytoplasmic before PDGF stimulation. After PDGF stimulation, localized at the cell membrane and peripheral region.
<b>Tissue Specificity</b>	Expressed in spleen, prostate, testis, uterus, small intestine and skeletal muscle. Among hematopoietic cell lines, expressed exclusively in B-cells. Not expressed in most tumor cell lines.
<b>Function</b>	function:Adapter protein for several members of the tyrosine kinase receptor family. Involved in multiple signaling pathways. May be involved in coupling from immunoreceptor to Ras signaling. Acts as a negative regulator of cytokine signaling in collaboration with CBL. Binds to EPOR and suppresses EPO-induced STAT5 activation, possibly through a masking effect on STAT5 docking sites in EPOR. Suppresses PDGF-induced mitogenesis. May induce cytoskeletal reorganization via interaction with VAV3.,PTM:Tyrosine phosphorylated by JAK2, KIT and other kinases activated by B-cell receptor in response to stimulation with cytokines, IL3, IL5, PDGF, IGF1, IGF2, CSF2/GM-CSF and cross-linking of the B-cell receptor complex.,similarity:Belongs to the SH2B adapter family.,similarity:Contains 1 PH domain.,similarity:Contains 1 SH2 domain.,subcellular location:Cytoplasmic before PDGF stimulation. After PDG



## Background

The protein encoded by this gene is expressed in B lymphocytes and contains pleckstrin homology and src homology 2 (SH2) domains. In Burkitt's lymphoma cell lines, it is tyrosine-phosphorylated in response to B cell receptor stimulation. Because it binds Shc independent of stimulation and Grb2 after stimulation, it appears to play a role in signal transduction from the receptor to the Shc/Grb2 pathway. [provided by RefSeq, Jun 2009],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western Blot analysis of HuvEc cells using APS Polyclonal Antibody