



Arkadia Polyclonal Antibody

Catalog No	YP-Ab-03712
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	RNF111
Protein Name	E3 ubiquitin-protein ligase Arkadia
Immunogen	The antiserum was produced against synthesized peptide derived from human RNF111. AA range:901-950
Specificity	Arkadia Polyclonal Antibody detects endogenous levels of Arkadia protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	RNF111; E3 ubiquitin-protein ligase Arkadia; RING finger protein 111
Observed Band	110kD
Cell Pathway	Nucleus . Cytoplasm . Nucleus, PML body . Upon TGF-beta treatment, translocates from nucleus to cytosol. .
Tissue Specificity	Broadly expressed.
Function	function:Acts in the NODAL pathway of mesoderm patterning during embryonic development. Acts downstream AXIN1 as an E3 ubiquitin-protein ligase which promotes the ubiquitination of inhibitory SMADs such as SMAD7, induces their proteasomal degradation and thereby enhances the transcriptional activity of TGF-beta and BMP. Activates Smad3/Smad4-dependent transcription by triggering signal-induced SnoN degradation.,pathway:Protein modification; protein ubiquitination.,similarity:Contains 1 RING-type zinc finger.,subcellular location:Upon TGF-beta treatment, translocates from nucleus to cytosol.,subunit:Interacts with SMAD6, SMAD7, AXIN1, AXIN2 and SKIL isoform SNON. Part of a complex containing RNF111, AXIN1 and SMAD7.,tissue specificity:Broadly expressed.,
Background	ring finger protein 111(RNF111) Homo sapiens The protein encoded by this gene is a nuclear RING-domain containing E3 ubiquitin ligase. This protein interacts with the transforming growth factor (TGF) -beta/NODAL signaling



pathway by promoting the ubiquitination and proteosomal degradation of negative regulators, like SMAD proteins, and thereby enhances TGF-beta target-gene transcription. As a modulator of the nodal signaling cascade, this gene plays a critical role in the induction of mesoderm during embryonic development. Alternative splicing of this gene results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2012],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of RNF111 Antibody. The lane on the right is blocked with the RNF111 peptide.



Western blot analysis of the lysates from Jurkat cells using RNF111 antibody.