


**(** Tel: 400-999-8863 ■ Email:Upingbio.163.com






## IFIT-5 Polyclonal Antibody

Catalog No	YP-Ab-03929
Catalog No	YP-AD-03929
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	IFIT5
Protein Name	Interferon-induced protein with tetratricopeptide repeats 5
Immunogen	The antiserum was produced against synthesized peptide derived from human IFIT5. AA range:281-330
Specificity	IFIT-5 Polyclonal Antibody detects endogenous levels of IFIT-5 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year  IFIT5; ISG58; RI58; Interferon-induced protein with tetratricopeptide repeats 5; IFIT-5; Interferon-induced 58 kDa protein; Retinoic acid- and interferon-inducible 58 kDa protein; P58
	IFIT5; ISG58; RI58; Interferon-induced protein with tetratricopeptide repeats 5; IFIT-5; Interferon-induced 58 kDa protein; Retinoic acid- and interferon-inducible
Synonyms	IFIT5; ISG58; RI58; Interferon-induced protein with tetratricopeptide repeats 5; IFIT-5; Interferon-induced 58 kDa protein; Retinoic acid- and interferon-inducible 58 kDa protein; P58
Synonyms Observed Band	IFIT5; ISG58; RI58; Interferon-induced protein with tetratricopeptide repeats 5; IFIT-5; Interferon-induced 58 kDa protein; Retinoic acid- and interferon-inducible 58 kDa protein; P58 50kD  Cell projection, ruffle membrane . Colocalized with DDX58/RIG-I at cell surface
Synonyms  Observed Band Cell Pathway	IFIT5; ISG58; RI58; Interferon-induced protein with tetratricopeptide repeats 5; IFIT-5; Interferon-induced 58 kDa protein; Retinoic acid- and interferon-inducible 58 kDa protein; P58 50kD  Cell projection, ruffle membrane . Colocalized with DDX58/RIG-I at cell surface ruffles. Localizes to actin-rich protrusions from the apical cell surface.
Synonyms  Observed Band  Cell Pathway  Tissue Specificity	IFIT5; ISG58; RI58; Interferon-induced protein with tetratricopeptide repeats 5; IFIT-5; Interferon-induced 58 kDa protein; Retinoic acid- and interferon-inducible 58 kDa protein; P58  50kD  Cell projection, ruffle membrane. Colocalized with DDX58/RIG-I at cell surface ruffles. Localizes to actin-rich protrusions from the apical cell surface.  Pancreas,Spleen,  induction:By interferons.,similarity:Belongs to the IFIT family.,similarity:Contains 8
Synonyms  Observed Band Cell Pathway Tissue Specificity Function	IFIT5; ISG58; RI58; Interferon-induced protein with tetratricopeptide repeats 5; IFIT-5; Interferon-induced 58 kDa protein; Retinoic acid- and interferon-inducible 58 kDa protein; P58 50kD  Cell projection, ruffle membrane. Colocalized with DDX58/RIG-I at cell surface ruffles. Localizes to actin-rich protrusions from the apical cell surface.  Pancreas,Spleen,  induction:By interferons.,similarity:Belongs to the IFIT family.,similarity:Contains 8 TPR repeats., induction:By interferons.,similarity:Belongs to the IFIT family.,similarity:Contains 8
Synonyms  Observed Band Cell Pathway Tissue Specificity Function Background matters needing	IFIT5; ISG58; RI58; Interferon-induced protein with tetratricopeptide repeats 5; IFIT-5; Interferon-induced 58 kDa protein; Retinoic acid- and interferon-inducible 58 kDa protein; P58  50kD  Cell projection, ruffle membrane. Colocalized with DDX58/RIG-I at cell surface ruffles. Localizes to actin-rich protrusions from the apical cell surface.  Pancreas,Spleen,  induction:By interferons.,similarity:Belongs to the IFIT family.,similarity:Contains 8 TPR repeats.,  induction:By interferons.,similarity:Belongs to the IFIT family.,similarity:Contains 8 TPR repeats.,



## **Products Images**



Western blot analysis of the lysates from HepG2 cells using IFIT5 antibody.