



MGST3 Polyclonal Antibody

Catalog No	YP-Ab-03957
Isotype	IgG
Reactivity	Human;Mouse
Applications	IHC;IF;ELISA
Gene Name	MGST3
Protein Name	Microsomal glutathione S-transferase 3
Immunogen	The antiserum was produced against synthesized peptide derived from human MGST3. AA range:8-57
Specificity	MGST3 Polyclonal Antibody detects endogenous levels of MGST3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MGST3; Microsomal glutathione S-transferase 3; Microsomal GST-3; Microsomal GST-III
Observed Band	
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Microsome membrane ; Multi-pass membrane protein . Microsome membrane ; Lipid-anchor . Membrane .
Tissue Specificity	Predominantly expressed in heart, skeletal muscle, and adrenal cortex. Also found in brain, placenta, liver, kidney, pancreas, thyroid, testis and ovary. Almost absent in lung, thymus and peripheral blood leukocytes.
Function	catalytic activity:RX + glutathione = HX + R-S-glutathione.,function:Also functions as a glutathione peroxidase.,similarity:Belongs to the MAPEG family.,tissue specificity:Predominantly expressed in heart, skeletal muscle, and adrenal cortex. Also found in brain, placenta, liver, kidney, pancreas, thyroid, testis and ovary. Almost absent in lung, thymus and peripheral blood leukocytes.,
Background	This gene encodes a member of the MAPEG (Membrane Associated Proteins in Eicosanoid and Glutathione metabolism) protein family. Members of this family are involved in the production of leukotrienes and prostaglandin E, important mediators of inflammation. This gene encodes an enzyme which catalyzes the conjugation of leukotriene A4 and reduced glutathione to produce leukotriene C4. This enzyme also demonstrates glutathione-dependent peroxidase activity



towards lipid hydroperoxides.[provided by RefSeq, May 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using MGST3 Antibody. The picture on the right is blocked with the synthesized peptide.