

Tel: 400-999-8863 
■ Email:Upingbio.163.com



## MRP-S17 Polyclonal Antibody

Catalog No	YP-Ab-04006
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;IHC;IF;ELISA
Gene Name	MRPS17
Protein Name	28S ribosomal protein S17 mitochondrial
lmmunogen	The antiserum was produced against synthesized peptide derived from human MRPS17. AA range:11-60
Specificity	MRP-S17 Polyclonal Antibody detects endogenous levels of MRP-S17 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/40000 IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MRPS17; RPMS17; HSPC011; 28S ribosomal protein S17; mitochondrial; MRP-S17; S17mt
Observed Band	22kD
Cell Pathway	Mitochondrion .
Tissue Specificity	Brain, Epithelium, Kidney, Umbilical cord blood, Uterus,
Function	similarity:Belongs to the ribosomal protein S17P family.,subunit:Component of the mitochondrial ribosome small subunit (28S) which comprises a 12S rRNA and about 30 distinct proteins.,
Background	Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein that belongs to the ribosomal protein \$17P family. The encoded protein is moderately

conserved between human mitochondrial and prokaryotic ribosomal proteins.



## UpingBio technology Co.,Ltd


📞 Tel: 400-999-8863 🗷 Emall:Upingbio.163.com




Pseudogenes corresponding to this gene are found

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.






Western blot analysis of lysates from NIH/3T3, RAW264.7, and A549 cells, using MRPS17 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HeLa cells using MRPS17 antibody.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).