




# AHNK Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-04887
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	IHC;IF
<b>Gene Name</b>	AHNAK PM227
<b>Protein Name</b>	Neuroblast differentiation-associated protein AHNAK (Desmoyokin)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 5720-5800
<b>Specificity</b>	AHNK Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	IHC-p 1:50-300. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	647kD
<b>Cell Pathway</b>	Nucleus.
<b>Tissue Specificity</b>	Cervix,Epithelium,Kidney,Liver,PCR rescued clones,Placenta,Tongue,Uterus,
<b>Function</b>	function:May be required for neuronal cell differentiation.,similarity:Contains 1 PDZ (DHR) domain.,subunit:Interacts with DYSF; the interaction is direct and Ca(2+)-independent.,
<b>Background</b>	function:May be required for neuronal cell differentiation.,similarity:Contains 1 PDZ (DHR) domain.,subunit:Interacts with DYSF; the interaction is direct and Ca(2+)-independent.,
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA, pH 9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200 (4° overnight). 3, Secondary antibody was diluted at 1:200 (room temperature, 45 min).