

**(** Tel: 400-999-8863 ■ Email:Upingbio.163.com





## MTF1 Polyclonal Antibody

| Catalog No                | YP-Ab-04990  |  |  |
|---------------------------|--|--|--|
| Isotype                   | IgG  |  |  |
| Reactivity                | Human;Mouse  |  |  |
| Applications              | WB;ELISA   |  |  |
| Gene Name                 | MTF1   |  |  |
| Protein Name              | Metal regulatory transcription factor 1 (MRE-binding transcription factor) (Transcription factor MTF-1)  |  |  |
| Immunogen                 | Synthesized peptide derived from human protein . at AA range: 260-340  |  |  |
| Specificity               | MTF1 Polyclonal Antibody detects endogenous levels of protein.   |  |  |
| Formulation               | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |  |  |
| Source                    | Polyclonal, Rabbit,IgG   |  |  |
| Purification              | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |  |  |
| Dilution                  | WB 1:500-2000 ELISA 1:5000-20000   |  |  |
| Concentration             | 1 mg/ml  |  |  |
| Purity                    | ≥90%   |  |  |
| Storage Stability         | -20°C/1 year   |  |  |
| Synonyms                  |  |  |  |
| Observed Band             | 82kD   |  |  |
| Cell Pathway              | Nucleus . Cytoplasm . Translocation to the nucleus is induced by metals  |  |  |
| Tissue Specificity        | Liver, Muscle,   |  |  |
| Function                  | function: Activates the metallothionein I promoter. Binds to the metal responsive  |  |  |
|                           | element (MRE).,similarity:Contains 6 C2H2-type zinc fingers.,  |  |  |
| Background                | This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], |  |  |
| matters needing attention | This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element  |  |  |
| matters needing           | This gene encodes a transcription factor that induces expression of metallothioneins and other genes involved in metal homeostasis in response to heavy metals such as cadmium, zinc, copper, and silver. The protein is a nucleocytoplasmic shuttling protein that accumulates in the nucleus upon heavy metal exposure and binds to promoters containing a metal-responsive element (MRE). [provided by RefSeq, Jul 2008], |  |  |



## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com



| Products Images |  |  |  |
|-----------------|--|--|--|
|                 |  |  |  |
|                 |  |  |  |
|                 |  |  |  |