




# MBIP1 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-05025
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	MBIP BM-015
<b>Protein Name</b>	MAP3K12-binding inhibitory protein 1 (MAPK upstream kinase-binding inhibitory protein) (MUK-binding inhibitory protein)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 1-80
<b>Specificity</b>	MBIP1 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	37kD
<b>Cell Pathway</b>	Nucleus. Cytoplasm. Shows a cytoplasmic localization when coexpressed with MAP3K12.
<b>Tissue Specificity</b>	Ubiquitous. High expression seen in the heart and lung.
<b>Function</b>	function:Inhibits the MAP3K12 activity to induce the activation of the JNK/SAPK pathway.,subcellular location:Shows a cytoplasmic localization when co-expressed with MAP3K12.,tissue specificity:Ubiquitous. High expression seen in the heart and lung.,
<b>Background</b>	function:Inhibits the MAP3K12 activity to induce the activation of the JNK/SAPK pathway.,subcellular location:Shows a cytoplasmic localization when co-expressed with MAP3K12.,tissue specificity:Ubiquitous. High expression seen in the heart and lung.,
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images



Western blot analysis of lysates from SH-SY5Y cells,  
primary antibody was diluted at 1:1000, 4° over night