



# MX1 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-05138
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	MX1
<b>Protein Name</b>	Interferon-induced GTP-binding protein Mx1 (Interferon-induced protein p78) (IFI-78K) (Interferon-regulated resistance GTP-binding protein MxA) (Myxoma resistance protein 1) (Myxovirus resistance prot
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 500-580
<b>Specificity</b>	MX1 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	72kD
<b>Cell Pathway</b>	Cytoplasm . Endoplasmic reticulum membrane ; Peripheral membrane protein ; Cytoplasmic side. Cytoplasm, perinuclear region . Binds preferentially to negatively charged phospholipids (PubMed:21900240). Colocalizes with CCHFV protein N in the perinuclear region (PubMed:15047845). .; [Isoform 2]: Cytoplasm . Nucleus . Translocates into the nuclei of HSV-1 infected cells (PubMed:20603636). .
<b>Tissue Specificity</b>	Primary B-Cells,Spleen,Uterus,
<b>Function</b>	function:May regulate the calcium channel activity of TRPCs. Shows activity against influenza virus and VSV, a rhabdovirus.,induction:By interferons.,similarity:Belongs to the dynamin family.,similarity:Contains 1 GED domain.,subunit:Interacts with TRPC1, TRPC3, TRPC4, TRPC5, TRPC6 and TRPC7.,
<b>Background</b>	This gene encodes a guanosine triphosphate (GTP)-metabolizing protein that participates in the cellular antiviral response. The encoded protein is induced by type I and type II interferons and antagonizes the replication process of several different RNA and DNA viruses. There is a related gene located adjacent to this gene on chromosome 21, and there are multiple pseudogenes located in a cluster

on chromosome 4. Alternative splicing results in multiple transcript variants.  
[provided by RefSeq, Sep 2013],

**matters needing  
attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For  
more information, please consult technical personnel.

## Products Images