

(Tel: 400-999-8863 ■ Email:Upingbio.163.com




GALM Polyclonal Antibody

Catalog No	YP-Ab-05304
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	GALM BLOCK25
Protein Name	Aldose 1-epimerase (EC 5.1.3.3) (Galactose mutarotase)
Immunogen	Synthesized peptide derived from human protein . at AA range: 40-120
Specificity	GALM Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Observed Band	37kD
Cell Pathway	37kD Cytoplasm .
Cell Pathway	Cytoplasm . Brain, Hematopoietic, Ovary, Skin, catalytic activity: Alpha-D-glucose = beta-D-glucose., function: Mutarotase converts alpha-aldose to the beta-anomer. It is active on D-glucose, L-arabinose, D-xylose, D-galactose, maltose and lactose., pathway: Carbohydrate metabolism; hexose
Cell Pathway Tissue Specificity	Cytoplasm . Brain, Hematopoietic, Ovary, Skin, catalytic activity: Alpha-D-glucose = beta-D-glucose., function: Mutarotase converts alpha-aldose to the beta-anomer. It is active on D-glucose, L-arabinose, D-xylose, D-galactose, maltose and lactose., pathway: Carbohydrate metabolism; hexose metabolism., similarity: Belongs to the aldose epimerase family., subunit: Monomer.,
Cell Pathway Tissue Specificity Function	Cytoplasm . Brain, Hematopoietic, Ovary, Skin, catalytic activity: Alpha-D-glucose = beta-D-glucose., function: Mutarotase converts alpha-aldose to the beta-anomer. It is active on D-glucose, L-arabinose, D-xylose, D-galactose, maltose and lactose., pathway: Carbohydrate metabolism; hexose metabolism., similarity: Belongs to the aldose epimerase family., subunit: Monomer., This gene encodes an enzyme that catalyzes the epimerization of hexose sugars such as glucose and galactose. The encoded protein is expressed in the cytoplasm and has a preference for galactose. The encoded protein may be required for normal galactose metabolism by maintaining the equilibrium of alpha
Cell Pathway Tissue Specificity Function Background matters needing	Cytoplasm . Brain, Hematopoietic, Ovary, Skin, catalytic activity: Alpha-D-glucose = beta-D-glucose., function: Mutarotase converts alpha-aldose to the beta-anomer. It is active on D-glucose, L-arabinose, D-xylose, D-galactose, maltose and lactose., pathway: Carbohydrate metabolism; hexose metabolism., similarity: Belongs to the aldose epimerase family., subunit: Monomer., This gene encodes an enzyme that catalyzes the epimerization of hexose sugars such as glucose and galactose. The encoded protein is expressed in the cytoplasm and has a preference for galactose. The encoded protein may be required for normal galactose metabolism by maintaining the equilibrium of alpha and beta anomers of galactose. [provided by RefSeq, Mar 2009],



Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night