



GCNT4 Polyclonal Antibody

Catalog No	YP-Ab-05372
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	GCNT4
Protein Name	Beta-1,3-galactosyl-O-glycosyl-glycoprotein beta-1,6-N-acetylglucosaminyltransferase 4 (EC 2.4.1.102) (Core 2-branching enzyme 3) (Core2-GlcNAc-transferase 3) (C2GnT3)
Immunogen	Synthesized peptide derived from human protein . at AA range: 60-140
Specificity	GCNT4 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	49kD
Cell Pathway	Golgi apparatus membrane ; Single-pass type II membrane protein .
Tissue Specificity	Predominantly expressed in thymus. Weakly expressed in pancreas, peripheral blood leukocytes, placenta, small intestine and stomach. Barely detectable in liver, spleen, lung and lymph node.
Function	catalytic activity:UDP-N-acetyl-D-glucosamine + beta-D-galactosyl-1,3-N-acetyl-D-galactosaminyl-R = UDP + beta-D-galactosyl-1,3-(N-acetyl-beta-D-glucosaminyl-1,6)-N-acetyl-D-galactosaminyl-R.,function:Glycosyltransferase that mediates core 2 O-glycan branching, an important step in mucin-type biosynthesis. Does not have core 4 O-glycan or I-branching enzyme activity.,online information:Core 2 beta-1,6-N-acetylglucosaminyltransferase 3.,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 14 family.,tissue specificity:Predominantly expressed in thymus. Weakly expressed in pancreas, peripheral blood leukocytes, placenta, small intestine and stomach. Barely detectable in liver, spleen, lung and lymph node.,



Background

catalytic activity:UDP-N-acetyl-D-glucosamine + beta-D-galactosyl-1,3-N-acetyl-D-galactosaminyl-R = UDP + beta-D-galactosyl-1,3-(N-acetyl-beta-D-glucosaminyl-1,6)-N-acetyl-D-galactosaminyl-R.,function:Glycosyltransferase that mediates core 2 O-glycan branching, an important step in mucin-type biosynthesis. Does not have core 4 O-glycan or I-branching enzyme activity.,online information:Core 2 beta-1,6-N-acetylglucosaminyltransferase 3.,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 14 family.,tissue specificity:Predominantly expressed in thymus. Weakly expressed in pancreas, peripheral blood leukocytes, placenta, small intestine and stomach. Barely detectable in liver, spleen, lung and lymph node.,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of Hela lysis, using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000