

Tel: 400-999-8863
 ■ Emall:Upingbio.163.com






B3GT5 Polyclonal Antibody

Catalog No YP-Ab-05373 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name B3GALT5 Protein Name Beta-1,3-galactosyltransferase 5 (Beta-1,3-GalTase 5) (IDP-Calbeta-GicNAc beta-1,3-galactosyltransferases 5) (IDP-Qalactose-beta-Hace) Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity B3GT5 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Solidate properties of the control of the contro		
Reactivity Human;Rat;Mouse; Applications WB;ELISA	Catalog No	YP-Ab-05373
Applications WB;ELISA Gene Name B3GALT5 Protein Name Beta-1,3-galactosyltransferase 5 (Beta-1,3-GalTase 5) (Beta3Gal-T5) (Beta3Gal-T5) (Beta3Gal-T5) (EC 2.4.1) (Beta-3-Gx-T5) (UDP-Gal:beta-GlcNAc beta-1,3-galactosyltransferase 5) (UDP-galactose:beta-N-ace Immunogen Synthesized peptide derived from human protein . at AA range: 50-130 Specificity B3GT5 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Golgi apparatus membrane ; Single-pass type II membrane protein . Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function: Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 0-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor, online information Beta-1,3-galactosyltransferase 5, online information: GlycoGene database, pathway. Protein modification; protein glycosylation, similarity Belongs to the glycosyltransferase 3 f family, tissue specificity Expressed in stomach, jejunum, colon, pancreas, small infestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes.	Isotype	lgG
Gene Name B3GALT5 Protein Name Beta-1,3-galactosyltransferase 5 (Beta-1,3-GalTase 5) (Beta3Gal-T5) (B2Gal-T5) (B2Ga	Reactivity	Human;Rat;Mouse;
Protein Name Beta-1,3-galactosyltransferase 5 (Beta-1,3-GalTase 5) (Beta3Gal-T5) (Beta3GalT5) (Beta-1,3-galactosyltransferase 5) (UDP-galactose:beta-N-ace	Applications	WB;ELISA
Beta3GalT5) (b3Gal-T5) (EC 2.4.1-) (Beta-3-Cx-T5) (UDP-Gal:beta-GicNAc beta-1,3-galactosyltransferase 5) (UDP-galictose:beta-N-ace Immunogen	Gene Name	B3GALT5
Specificity B3GT5 Polyclonal Antibody detects endogenous levels of protein.	Protein Name	Beta-1,3-galactosyltransferase 5 (Beta-1,3-GalTase 5) (Beta3Gal-T5) (Beta3GalT5) (b3Gal-T5) (EC 2.4.1) (Beta-3-Gx-T5) (UDP-Gal:beta-GlcNAc beta-1,3-galactosyltransferase 5) (UDP-galactose:beta-N-ace
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Golgi apparatus membrane; Single-pass type II membrane protein. Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc/beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor, online information:Beta-1.3-galactosyltransferase 5, online information:GlycoGene database, pathway:Protein modification; protein glycosylation, similarity;Belongs to the glycosyltransferase 31 family, tissue specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancer; Alternatively spliced transcript variants	Immunogen	Synthesized peptide derived from human protein . at AA range: 50-130
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Golgi apparatus membrane; Single-pass type II membrane protein . Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor, online information:GlycoGene database, pathway: Protein modification: protein glycosylation, similarity. Belongs to the glycosyltransferase 3.1 family, tissue specificity: Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancer. Alternatively spliced transcript variants	Specificity	B3GT5 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Golgi apparatus membrane; Single-pass type II membrane protein. Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor, online information:Beta-1,3-galactosyltransferase 5, online information:GlycoGene database, pathway:Protein modification; protein glycosylation, similarity:Belongs to the glycosyltransferase 31 family, tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes., Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Golgi apparatus membrane; Single-pass type II membrane protein. Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor, online information:Beta-1,3-galactosyltransferase 5,online information:GlycoGene database,pathway:Protein modification; protein glycosyltransinlarity:Belongs to the glycosyltransferase 31 family_tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Golgi apparatus membrane; Single-pass type II membrane protein . Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor.,online information:GlycoGene database, pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 3 family.,tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Purification	•
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Golgi apparatus membrane; Single-pass type II membrane protein. Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor, online information:Beta-1,3-galactosyltransferase 5, online information:GlycoGene database, pathway:Protein modification; protein glycosylation., similarity:Belongs to the glycosyltransferase 31 family, tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes., Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Synonyms Observed Band 34kD Cell Pathway Golgi apparatus membrane; Single-pass type II membrane protein. Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function: Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor.,online information:Beta-1,3-galactosyltransferase 5,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 31 family.,tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes., Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Concentration	1 mg/ml
Observed Band Golgi apparatus membrane; Single-pass type II membrane protein. Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor.,online information:Beta-1,3-galactosyltransferase 5,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 31 family., tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes., Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Purity	≥90%
Observed Band 34kD Cell Pathway Golgi apparatus membrane; Single-pass type II membrane protein. Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor.,online information:BlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 31 family.,tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes., Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Storage Stability	-20°C/1 year
Cell Pathway Golgi apparatus membrane; Single-pass type II membrane protein. Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor.,online information:Beta-1,3-galactosyltransferase 5,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 31 family.,tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes., Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Synonyms	
Tissue Specificity Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor.,online information:Beta-1,3-galactosyltransferase 5,online information:GlycoGene database, pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 31 family.,tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes., This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Observed Band	34kD
gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes. Function function:Catalyzes the transfer of Gal to GlcNAc-based acceptors with a preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor.,online information:Beta-1,3-galactosyltransferase 5,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 31 family.,tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes., This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Cell Pathway	Golgi apparatus membrane ; Single-pass type II membrane protein .
preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor.,online information:Beta-1,3-galactosyltransferase 5,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 31 family.,tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and peripheral blood leukocytes., Background This gene encodes a member of a family of membrane-bound glycoproteins. The encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Tissue Specificity	gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver,
encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants	Function	preference for the core3 O-linked glycan GlcNAc(beta1,3)GalNAc structure. Can use glycolipid LC3Cer as an efficient acceptor.,online information:Beta-1,3-galactosyltransferase 5,online information:GlycoGene database,pathway:Protein modification; protein glycosylation.,similarity:Belongs to the glycosyltransferase 31 family.,tissue specificity:Expressed in stomach, jejunum, colon, pancreas, small intestine, testis and gastrointestinal and pancreatic cancer cell lines. Hardly detected in lung, liver, adrenal gland and
	Background	encoded protein may synthesize type 1 Lewis antigens, which are elevated in gastrointestinal and pancreatic cancers. Alternatively spliced transcript variants



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



variants has not been determined. [provided by RefSeq, Jul 2013],

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



