



RPA1 Polyclonal Antibody

| | |
|--------------------|--|
| Catalog No | YP-Ab-05529 |
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB;ELISA |
| Gene Name | POLR1A |
| Protein Name | DNA-directed RNA polymerase I subunit RPA1 (RNA polymerase I subunit A1) (EC 2.7.7.6) (A190) (DNA-directed RNA polymerase I largest subunit) (DNA-directed RNA polymerase I subunit A) (RNA polymerase I |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | RPA1 Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 189kD |
| Cell Pathway | Nucleus, nucleolus . Chromosome . |
| Tissue Specificity | Colon,Skin,Uterus, |
| Function | catalytic activity:Nucleoside triphosphate + RNA(n) = diphosphate + RNA(n+1).,function:DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Largest and catalytic core component of RNA polymerase I which synthesizes ribosomal RNA precursors. Forms the polymerase active center together with the second largest subunit. A single stranded DNA template strand of the promoter is positioned within the central active site cleft of Pol I. A bridging helix emanates from RPA1 and crosses the cleft near the catalytic site and is thought to promote translocation of Pol I by acting as a ratchet that moves the RNA-DNA hybrid through the active site by switching from straight to bent conformations at each step of nucleotide addition.,PTM:Phosphorylated.,similarity:Belongs to the RNA polymerase beta' chain family.,subunit:Compo |
| Background | The protein encoded by this gene is the largest subunit of the RNA polymerase I complex. The encoded protein represents the catalytic subunit of the complex, |

which transcribes DNA into ribosomal RNA precursors. Defects in this gene are a cause of the Cincinnati type of acrofacial dysostosis. [provided by RefSeq, May 2016],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images