

(Tel: 400-999-8863 ■ Email:Upingbio.163.com




DIRA2 Polyclonal Antibody

Catalog No	YP-Ab-05616
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	DIRAS2
Protein Name	GTP-binding protein Di-Ras2 (Distinct subgroup of the Ras family member 2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DIRA2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year 21kD
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year 21kD Cell membrane ; Lipid-anchor ; Cytoplasmic side .
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year 21kD Cell membrane; Lipid-anchor; Cytoplasmic side. Highly expressed in brain. function:Displays low GTPase activity and exist predominantly in the GTP-bound form.,similarity:Belongs to the small GTPase superfamily. Di-Ras family.,tissue specificity:Highly expressed in brain., DIRAS2 belongs to a distinct branch of the functionally diverse Ras (see HRAS;
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year 21kD Cell membrane; Lipid-anchor; Cytoplasmic side. Highly expressed in brain. function:Displays low GTPase activity and exist predominantly in the GTP-bound form.,similarity:Belongs to the small GTPase superfamily. Di-Ras family.,tissue specificity:Highly expressed in brain.,
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year 21kD Cell membrane; Lipid-anchor; Cytoplasmic side. Highly expressed in brain. function:Displays low GTPase activity and exist predominantly in the GTP-bound form.,similarity:Belongs to the small GTPase superfamily. Di-Ras family.,tissue specificity:Highly expressed in brain., DIRAS2 belongs to a distinct branch of the functionally diverse Ras (see HRAS; MIM 190020) superfamily of monomeric GTPases.[supplied by OMIM, Apr 2004],



Products Images



Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night