

(Tel: 400-999-8863 ■ Email:Upingbio.163.com





HXA9 Polyclonal Antibody

Catalog No	YP-Ab-05649
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	HOXA9 HOX1G
Protein Name	Homeobox protein Hox-A9 (Homeobox protein Hox-1G)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	HXA9 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	29kD
Cell Pathway	Nucleus.
Tissue Specificity	Bone marrow,Colon,
Function	disease:A chromosomal aberration involving HOXA9 is found in a form of acute myeloid leukemia. Translocation t(7;11)(p15;p15) with NUP98.,disease:A chromosomal aberration involving HOXA9 may contribute to disease progression in chronic myeloid leukemia. Translocation t(7;17)(p15;q23) with MSI2.,function:Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis.,similarity:Belongs to the Abd-B homeobox family.,similarity:Contains 1 homeobox DNA-binding domain.,
Background	In vertebrates, the genes encoding the class of transcription factors called homeobox genes are found in clusters named A, B, C, and D on four separate chromosomes. Expression of these proteins is spatially and temporally regulated during embryonic development. This gene is part of the A cluster on chromosome 7 and encodes a DNA-binding transcription factor which may regulate gene expression, morphogenesis, and differentiation. This gene is highly similar to the abdominal-B (Abd-B) gene of Drosophila. A specific translocation event which



UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 💌 Emall:Upingbio.163.com



causes a fusion between this gene and the NUP98 gene has been associated with myeloid leukemogenesis. Read-through transcription exists between this gene and the upstream homeobox A10 (HOXA10) gene.[provided by RefSeq, Mar 2011],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

