



HMCS1 Polyclonal Antibody

Catalog No	YP-Ab-05674
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	HMGCS1 HMGCS
Protein Name	Hydroxymethylglutaryl-CoA synthase, cytoplasmic (HMG-CoA synthase) (EC 2.3.3.10) (3-hydroxy-3-methylglutaryl coenzyme A synthase)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	HMCS1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	57kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Brain,Epithelium,Fetal adrenal gland,Fibroblast,Liver,Lung,
Function	catalytic activity:Acetyl-CoA + H(2)O + acetoacetyl-CoA = (S)-3-hydroxy-3-methylglutaryl-CoA + CoA.,function:This enzyme condenses acetyl-CoA with acetoacetyl-CoA to form HMG-CoA, which is the substrate for HMG-CoA reductase.,pathway:Metabolic intermediate biosynthesis; mevalonic acid biosynthesis; (R)-mevalonic acid from acetyl-CoA: step 2/3.,similarity:Belongs to the HMG-CoA synthase family.,
Background	catalytic activity:Acetyl-CoA + H(2)O + acetoacetyl-CoA = (S)-3-hydroxy-3-methylglutaryl-CoA + CoA.,function:This enzyme condenses acetyl-CoA with acetoacetyl-CoA to form HMG-CoA, which is the substrate for HMG-CoA reductase.,pathway:Metabolic intermediate biosynthesis; mevalonic acid biosynthesis; (R)-mevalonic acid from acetyl-CoA: step 2/3.,similarity:Belongs to the HMG-CoA synthase family.,
matters needing attention	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images