



# A4GAT Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-05688
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	A4GALT A14GALT A4GALT1
<b>Protein Name</b>	Lactosylceramide 4-alpha-galactosyltransferase (EC 2.4.1.228) (Alpha-1,4-N-acetylglucosaminyltransferase) (Alpha-1,4-galactosyltransferase) (Alpha4Gal-T1) (CD77 synthase) (Globotriaosylceramide syntha
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	A4GAT Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	38kD
<b>Cell Pathway</b>	Golgi apparatus membrane ; Single-pass type II membrane protein .
<b>Tissue Specificity</b>	Ubiquitous. Highly expressed in kidney, heart, spleen, liver, testis and placenta.
<b>Function</b>	catalytic activity:UDP-galactose + beta-D-galactosyl-(1->4)-D-glucosyl-(1<->1)-ceramide = UDP + alpha-D-galactosyl-(1->4)-beta-D-galactosyl-(1->4)-D-glucosyl-(1<->1)-ceramide. ,domain:The conserved DXD motif is involved in enzyme activity.,function:Necessary for the biosynthesis of the Pk antigen of blood histogroup P. Catalyzes the transfer of galactose to lactosylceramide and galactosylceramide. Necessary for the synthesis of the receptor for bacterial verotoxins.,online information:GlycoGene database,online information:Lactosylceramide 4-alpha-galactosyltransferase,pathway:Protein modification; protein glycosylation.,polymorphism:Different combinations or absence of the P blood group system antigens define 5 different phenotypes: P1, P2, P1(k), P2(k), and p. Genetic variation in A4GALT determines the p phenotype, which is rare and does not express any antigens. It is also known as null



## Background

alpha 1,4-galactosyltransferase(A4GALT) Homo sapiens The protein encoded by this gene catalyzes the transfer of galactose to lactosylceramide to form globotriaosylceramide, which has been identified as the P(k) antigen of the P blood group system. This protein, a type II membrane protein found in the Golgi, is also required for the synthesis of the bacterial verotoxins receptor. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2015],


## matters needing attention

Avoid repeated freezing and thawing!

## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night