

(Tel: 400-999-8863 ■ Email:Upingbio.163.com



LHX8 Polyclonal Antibody

This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with		
Reactivity Human;Mouse Applications WB;ELISA Gene Name LHX8 Protein Name LIM/homeobox protein Lhx8 (LIM homeobox protein 8) Immunogen Synthesized peptide derived from human protein . at AA range: 150-230 Specificity LHX8 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus . Tissue Specificity Peripheral blood, Testis, Function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells, similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: conta	Catalog No	YP-Ab-05693
Applications WB;ELISA Gene Name LHX8 Protein Name LIM/homeobox protein Lhx8 (LIM homeobox protein 8) Immunogen Synthesized peptide derived from human protein . at AA range: 150-230 Specificity LHX8 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus . Tissue Specificity Peripheral blood, Testis, Function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells, similarity-Contains 1 homeobox DNA-binding domain, similarity-Contains 1 LIM zinc-binding domain,, similarity-Contains 1 LIM zinc-binding domain,, similarity-Contains 1 The protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LiM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odonoma formation. Alternative splicing of this gene results in multiple transcription donomain formation. Alternative splicing of this gene results in multiple transcription donomain.	Isotype	IgG
Gene Name LHX8 Protein Name LIM/homeobox protein Lhx8 (LIM homeobox protein 8) Immunogen Synthesized peptide derived from human protein . at AA range: 150-230 Specificity LHX8 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus . Tissue Specificity Peripheral blood, Testis, Function function: Transcription factor involved in differentiation of certain neurons and messenchymal cells, similarity: Contains 1 homeobox DNA-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 1 homeobox planting of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-b	Reactivity	Human;Mouse
Immunogen Synthesized peptide derived from human protein . at AA range: 150-230	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 150-230 Specificity LHX8 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus . Function function:Transcription factor involved in differentiation of certain neurons and mesenchymal cells, similarity:Contains 1 homeobox DNA-binding domain, similarity:Contains 2 LIM zinc-binding domains, similarity:Contains 2 LIM zinc-binding domains, similarity:Contains 2 LIM zinc-binding domains, similarity:The protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated wi	Gene Name	LHX8
Specificity LHX8 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus. Tissue Specificity Peripheral blood, Testis, Function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells. similarity: Contains 1 homeobox DNA-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domains. The protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc linger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ougenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odonloma formation. Alternative splicing of this gene results in multiply transcript	Protein Name	LIM/homeobox protein Lhx8 (LIM homeobox protein 8)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus . Tissue Specificity Peripheral blood,Testis, Function function:Transcription factor involved in differentiation of certain neurons and mesenchymal cells, similarity:Contains 1 homeobox DNA-binding domain., similarity:Contains 1 LIM zinc-binding domain., similarity:Contains 2 LIM zinc-binding domain., similarity:These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This gene is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Immunogen	Synthesized peptide derived from human protein . at AA range: 150-230
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus . Tissue Specificity Peripheral blood,Testis, Function function:Transcription factor involved in differentiation of certain neurons and mesenchymal cells., similarity:Contains 1 homeobox DNA-binding domain., similarity:Contains 2 LIM zinc-binding domain., similarity:Contains 2 LIM zinc-binding domain., similarity:Contains 2 LIM zinc-binding domain, similarity:Contains 2 LIM zinc-binding domain, and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcription factor that plays a role in tooth	Specificity	LHX8 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus Tissue Specificity Peripheral blood, Testis, function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells, similarity: Contains 1 homeobox DNA-binding domain, similarity: Contains 1 LIM zinc-binding domain, similarity: Contains 2 LIM zinc-binding domain, similarity: Synonymia and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ogenesis and in neuronal differentiation. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ogenesis and in neuronal differentiation. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ogenesis and in neuronal differentiation. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ogenesis and in neuronal differentiation. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ogenesis and in neuronal differentiation.	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus . Tissue Specificity Peripheral blood, Testis, Function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells., similarity: Contains 1 homeobox DNA-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 1 tulm zinc-binding domain., similarity: Contains 1 tulm zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 3 LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 39kD Cell Pathway Nucleus . Tissue Specificity Peripheral blood,Testis, Function function:Transcription factor involved in differentiation of certain neurons and mesenchymal cells.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 1 LIM zinc-binding domain.,similarity:Contains 2 LIM zinc-binding domains., in the protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 39kD Cell Pathway Nucleus . Tissue Specificity Peripheral blood,Testis, Function function:Transcription factor involved in differentiation of certain neurons and mesenchymal cells.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 1 LIM zinc-binding domain.,similarity:Contains 2 LIM zinc-binding domains., the protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Synonyms Observed Band 39kD Cell Pathway Nucleus . Function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells., similarity: Contains 1 homeobox DNA-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 3 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 3 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 3 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 3 LIM zinc-binding domain., similarity: Contains 3 LIM zinc-binding domain., similarity: Contains 3 LIM zinc-binding and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Concentration	1 mg/ml
Observed Band Gell Pathway Nucleus Tissue Specificity Peripheral blood, Testis, Function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells., similarity: Contains 1 homeobox DNA-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domains., Background The protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in ogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Purity	≥90%
Observed Band Cell Pathway Nucleus . Peripheral blood, Testis, function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells., similarity: Contains 1 homeobox DNA-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domains., Background The protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Storage Stability	-20°C/1 year
Tissue Specificity Peripheral blood, Testis, function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells., similarity: Contains 1 homeobox DNA-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domains., Background The protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Synonyms	
Tissue Specificity Peripheral blood, Testis, function function: Transcription factor involved in differentiation of certain neurons and mesenchymal cells., similarity: Contains 1 homeobox DNA-binding domain., similarity: Contains 1 LIM zinc-binding domain., similarity: Contains 2 LIM zinc-binding domains., Background The protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Observed Band	39kD
Function function:Transcription factor involved in differentiation of certain neurons and mesenchymal cells.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 2 LIM zinc-binding domain.,similarity:Contains 2 LIM zinc-binding domain.,similarity:Contains 2 LIM zinc-binding domains., The protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Cell Pathway	Nucleus .
mesenchymal cells.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 2 LIM zinc-binding domain.,similarity:Contains 2 LIM zinc-binding domains., Background The protein encoded by this gene is a member of the LIM homeobox family of proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Tissue Specificity	Peripheral blood,Testis,
proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript	Function	mesenchymal cells.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 1 LIM zinc-binding domain.,similarity:Contains 2 LIM
	Background	proteins, which are involved in patterning and differentiation of various tissue types. These proteins contain two tandemly repeated cysteine-rich double-zinc finger motifs known as LIM domains, in addition to a DNA-binding homeodomain. This family member is a transcription factor that plays a role in tooth morphogenesis. It is also involved in oogenesis and in neuronal differentiation. This gene is a candidate gene for cleft palate, and it is also associated with odontoma formation. Alternative splicing of this gene results in multiple transcript



UpingBio technology Co.,Ltd

Tel: 400-999-8863
 ■ Email:Upingbio.163.com




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night