



MY18A Polyclonal Antibody

Catalog No	YP-Ab-05785
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	MYO18A KIAA0216 MYSPDZ
Protein Name	Unconventional myosin-XVIIIa (Molecule associated with JAK3 N-terminus) (MAJN) (Myosin containing a PDZ domain)
Immunogen	Synthesized peptide derived from human protein . at AA range: 1920-2000
Specificity	MY18A Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	225kD
Cell Pathway	Golgi apparatus . Golgi apparatus, trans-Golgi network . Golgi outpost . Cytoplasm, cytoskeleton, microtubule organizing center . Recruited to the Golgi apparatus by GOLPH3 (PubMed:19837035). Localizes to the postsynaptic Golgi apparatus region, also named Golgi outpost, which shapes dendrite morphology by functioning as sites of acentrosomal microtubule nucleation (By similarity). . ; [Isoform 1]: Endoplasmic reticulum-Golgi intermediate compartment . Cytoplasm, cytoskeleton . Colocalizes with actin. . ; [Isoform 2]: Cytoplasm . Lacks the PDZ domain (PubMed:12761286). Diffusely localized in the cytoplasm (PubMed:12761286). . ; [Isoform 5]: Cell surface .
Tissue Specificity	Amygdala,Brain,Epithelium,Liver,Peripheral blood,Platelet,Testis,
Function	caution:The TIAF1 protein is coded in the 3'-UTR region of MYO18A.,function:May be involved in the maintenance of the stromal cell architectures required for cell to cell contact.,similarity:Contains 1 IQ domain.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 PDZ (DHR) domain.,subcellular location:Colocalizes with actin.,subcellular location:Lacks the PDZ domain. Diffusely localized in the cytoplasm.,subunit:Homodimer. Interacts with JAK3 and actin.,

Background

caution:The TIAF1 protein is coded in the 3'-UTR region of MYO18A.,function:May be involved in the maintenance of the stromal cell architectures required for cell to cell contact.,similarity:Contains 1 IQ domain.,similarity:Contains 1 myosin head-like domain.,similarity:Contains 1 PDZ (DHR) domain.,subcellular location:Colocalizes with actin.,subcellular location:Lacks the PDZ domain. Diffusely localized in the cytoplasm.,subunit:Homodimer. Interacts with JAK3 and actin.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images