

(Tel: 400-999-8863 ■ Email:Upingbio.163.com



RAB1A Polyclonal Antibody

| Catalog No | YP-Ab-06040 |
|---------------------------|--|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse |
| Applications | WB;ELISA |
| Gene Name | RAB1A RAB1 |
| Protein Name | Ras-related protein Rab-1A (YPT1-related protein) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 50-130 |
| Specificity | RAB1A Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 22kD |
| Cell Pathway | Golgi apparatus . Endoplasmic reticulum . Early endosome . Cytoplasm, cytosol . Membrane . Melanosome . Alternates between membrane-associated and cytosolic forms |
| Tissue Specificity | Brain, Colon, Placenta, Platelet, Tongue, |
| Function | function:Probably required for transit of protein from the ER through Golgi compartment. Binds GTP and GDP and possesses intrinsic GTPase activity.,PTM:Phosphorylated by CDC2 kinase during mitosis.,similarity:Belongs to the small GTPase superfamily. Rab family.,subunit:May interact with YIPF5., |
| Background | This gene encodes a member of the Ras superfamily of GTPases. Members of the gene family cycle between inactive GDP-bound and active GTP-bound forms. This small GTPase controls vesicle traffic from the endoplasmic reticulum to the Golgi apparatus. Multiple alternatively spliced transcript variants have been identified for this gene which encode different protein isoforms. [provided by RefSeq, Oct 2008], |
| matters needing attention | Avoid repeated freezing and thawing! |
| | |



UpingBio technology Co.,Ltd






Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night