

**(** Tel: 400-999-8863 ■ Email:Upingbio.163.com





## SCN1B Polyclonal Antibody

| Catalog No         | YP-Ab-06203   |
|--------------------|---|
| Isotype            | IgG   |
| Reactivity         | Human;Rat;Mouse   |
| Applications       | WB;ELISA  |
| Gene Name          | SCN1B   |
| Protein Name       | Sodium channel subunit beta-1   |
| Immunogen          | Synthesized peptide derived from part region of human protein   |
| Specificity        | SCN1B Polyclonal Antibody detects endogenous levels of protein.   |
| Formulation        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  |
| Source             | Polyclonal, Rabbit,IgG  |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution           | WB 1:500-2000 ELISA 1:5000-20000  |
| Concentration      | 1 mg/ml   |
| Purity             | ≥90%  |
| Storage Stability  | -20°C/1 year  |
| Synonyms           |   |
| Observed Band      | 23kD  |
| Cell Pathway       | [Isoform 1]: Cell membrane; Single-pass type I membrane protein. Perikaryon. Cell projection. Cell projection, axon. Detected at nodes of Ranvier on the sciatic nerve; [Isoform 2]: Perikaryon. Cell projection. Secreted. Detected on Purkinje cells and their cell projections and on neuronal cell projections.   |
| Tissue Specificity | The overall expression of isoform 1 and isoform 2 is very similar. Isoform 1 is abundantly expressed in skeletal muscle, heart and brain. Isoform 2 is highly expressed in brain and skeletal muscle and present at a very low level in heart, placenta, lung, liver, kidney and pancreas. In brain, isoform 2 is most abundant in the cerebellum, followed by the cerebral cortex and occipital lobe, while isoform 1 levels are higher in the cortex compared to the cerebellum. Isoform 2 is expressed in many regions of the brain, including cerebellar Purkinje cells, cortex pyramidal neurons and many of the neuronal fibers throughout the brain (at protein level). Also detected in dorsal root ganglion, in fibers of the spinal nerve and in cortical neurons and their processes (at protein level). |
| Function           | disease:Defects in SCN1B are the cause of generalized epilepsy with febrile seizures plus type 1 (GEFS+1) [MIM:604233]. Generalized epilepsy with febrile seizures-plus refers to a rare autosomal dominant, familial condition with incomplete penetrance and large intrafamilial variability. Patients display febrile seizures persisting sometimes beyond the age of 6 years and/or a variety of  |



**Background** 


matters needing

**Usage suggestions** 

attention

## UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 🗷 Emall:Upingbio.163.com




| gen<br>part<br>ass<br>the<br>isof             | eralize<br>tial sei<br>embly<br>sodiur<br>orms f           | ed seizú<br>zures, v<br>, expres<br>n chanr<br>rom bra   | res ofter<br>vith a va<br>sion, an<br>el. The<br>in, skele           | n precipi<br>riable de<br>d functio<br>subunit                       | itated by<br>egree of<br>onal mo<br>beta-1 o<br>cle, and       | y fever a<br>severity<br>dulation<br>can mod<br>heart. I                | at age 6<br>y.,function<br>of the hold                                   | orile seizu<br>years or r<br>on:Crucial<br>aeterotrime<br>ultiple alpl<br>ciation with                     | nore, ar<br>in the<br>eric com<br>na subu                       | nplex of                             |
|---|--|--|--|--|--|---|--|--|---|--------------------------------------|
| gen<br>The<br>provinac<br>this<br>sync<br>enc | eration<br>y are o<br>vides o<br>ctivatio<br>gene<br>drome | n and procompose thannel notes that the contract of the contra | opagation<br>ed of on<br>activity of<br>gene en<br>genera<br>defects | on of act<br>e alpha<br>and the<br>codes a<br>lized epi<br>in cardia | tion pote<br>and two<br>beta-1 s<br>sodium<br>ilepsy wac condi | entials ir<br>beta su<br>subunit r<br>channe<br>ith febril<br>uction. N | n muscle<br>ubunits,<br>nodulate<br>el beta-1<br>le seizu<br>//ultiple f | s that fund<br>e and neur<br>where the<br>es the kind<br>subunit.<br>res plus, E<br>transcript ve.[provide | ronal ce<br>alpha<br>etics of<br>Mutatio<br>Brugada<br>variants | ells.<br>subunit<br>channel<br>ns in |
| Avo   | id repe  | eated fr   | eezing a   | ınd thaw   | ring!  |   |  |  |   |                                      |

This product can be used in immunological reaction related experiments. For

## **Products Images**

more information, please consult technical personnel.



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night