



# SYN3 Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-06251
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	SYN3
<b>Protein Name</b>	Synapsin-3 (Synapsin III)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein. AA range 530-580
<b>Specificity</b>	SYN3 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	63kD
<b>Cell Pathway</b>	Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Peripheral membrane protein localized to the cytoplasmic surface of synaptic vesicles.
<b>Tissue Specificity</b>	Neuron specific. Detected predominantly in brain.
<b>Function</b>	function:May be involved in the regulation of neurotransmitter release and synaptogenesis.,miscellaneous:Regulated by calcium. Calcium inhibits ATP binding to the C-domain.,similarity:Belongs to the synapsin family.,subcellular location:Peripheral membrane protein localized to the cytoplasmic surface of synaptic vesicles.,subunit:Interacts with CAPON.,tissue specificity:Neuron specific. Detected predominantly in brain.,
<b>Background</b>	This gene is a member of the synapsin gene family. Synapsins encode neuronal phosphoproteins which associate with the cytoplasmic surface of synaptic vesicles. Family members are characterized by common protein domains, and they are implicated in synaptogenesis and the modulation of neurotransmitter release, suggesting a potential role in several neuropsychiatric diseases. The protein encoded by this gene shares the synapsin family domain model, with domains A, C, and E exhibiting the highest degree of conservation. The protein

contains a unique domain J, located between domains C and E. Based on this gene's localization to 22q12.3, a possible schizophrenia susceptibility locus, and the established neurobiological roles of the synapsins, this family member may represent a candidate gene for schizophrenia. The TIMP3 gene is located within an intron of this gene and is transcribed in the opposite direction.

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images