



# UBP37 Polyclonal Antibody

|                                  |  |
|----------------------------------|--|
| <b>Catalog No</b>                | YP-Ab-06344  |
| <b>Isotype</b>                   | IgG  |
| <b>Reactivity</b>                | Human;Rat;Mouse;   |
| <b>Applications</b>              | WB;ELISA   |
| <b>Gene Name</b>                 | USP37 KIAA1594   |
| <b>Protein Name</b>              | Ubiquitin carboxyl-terminal hydrolase 37 (EC 3.4.19.12) (Deubiquitinating enzyme 37) (Ubiquitin thioesterase 37) (Ubiquitin-specific-processing protease 37)   |
| <b>Immunogen</b>                 | Synthesized peptide derived from part region of human protein  |
| <b>Specificity</b>               | UBP37 Polyclonal Antibody detects endogenous levels of protein.  |
| <b>Formulation</b>               | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |
| <b>Source</b>                    | Polyclonal, Rabbit,IgG   |
| <b>Purification</b>              | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| <b>Dilution</b>                  | WB 1:500-2000 ELISA 1:5000-20000   |
| <b>Concentration</b>             | 1 mg/ml  |
| <b>Purity</b>                    | ≥90%   |
| <b>Storage Stability</b>         | -20°C/1 year   |
| <b>Synonyms</b>                  |  |
| <b>Observed Band</b>             | 107kD  |
| <b>Cell Pathway</b>              | nucleus,   |
| <b>Tissue Specificity</b>        | Expressed in brain and prostate.   |
| <b>Function</b>                  | catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 3 UIM (ubiquitin-interacting motif) repeats.,tissue specificity:Expressed in brain and prostate., |
| <b>Background</b>                | catalytic activity:Ubiquitin C-terminal thioester + H(2)O = ubiquitin + a thiol.,similarity:Belongs to the peptidase C19 family.,similarity:Contains 3 UIM (ubiquitin-interacting motif) repeats.,tissue specificity:Expressed in brain and prostate., |
| <b>matters needing attention</b> | Avoid repeated freezing and thawing!   |
| <b>Usage suggestions</b>         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |



## Products Images