

**(** Tel: 400-999-8863 ■ Email:Upingbio.163.com





## **CCNY Polyclonal Antibody**

Catalog No	YP-Ab-06436
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	CCNY C10orf9 CBCP1 CFP1
Protein Name	Cyclin-Y (Cyc-Y) (Cyclin box protein 1) (Cyclin fold protein 1) (cyclin-X)
Immunogen	Synthesized peptide derived from human protein . at AA range: 270-350
Specificity	CCNY Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  37kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90%  -20°C/1 year  37kD  Cell membrane ; Lipid-anchor ; Cytoplasmic side .; [Isoform 3]: Nucleus.
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  37kD  Cell membrane ; Lipid-anchor ; Cytoplasmic side .; [Isoform 3]: Nucleus.  Widely expressed.  similarity:Belongs to the cyclin family. Cyclin Y subfamily.,similarity:Contains 1
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  37kD  Cell membrane; Lipid-anchor; Cytoplasmic side .; [Isoform 3]: Nucleus.  Widely expressed.  similarity:Belongs to the cyclin family. Cyclin Y subfamily.,similarity:Contains 1 cyclin N-terminal domain.,  Cyclins, such as CCNY, control cell division cycles and regulate
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  37kD  Cell membrane; Lipid-anchor; Cytoplasmic side .; [Isoform 3]: Nucleus.  Widely expressed.  similarity:Belongs to the cyclin family. Cyclin Y subfamily.,similarity:Contains 1 cyclin N-terminal domain.,  Cyclins, such as CCNY, control cell division cycles and regulate cyclin-dependent kinases (e.g., CDC2; MIM 116940) (Li et al., 2009 [PubMed 18060517]).[supplied by OMIM, May 2009],




## UpingBio technology Co.,Ltd

**(** Tel: 400-999-8863 ■ Email:Upingbio.163.com



## **Products Images**



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night