

(Tel: 400-999-8863 ■ Email:Upingbio.163.com



ZN750 Polyclonal Antibody

Catalog No	YP-Ab-06495
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	ZNF750
Protein Name	Zinc finger protein 750
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	ZN750 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	79kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in the skin, prostate, lung, placenta and thymus, and at low level in T-cells. Not expressed in peripheral blood leukocytes, pancreas and brain. Clearly expressed in primary keratinocytes but not in fibroblasts.
Function	caution:Despite its name, it does not contain a canonical C2H2-type zinc-finger.,disease:Defects in ZNF750 are the cause of Seborrhea-like dermatitis with psoriasiform [MIM:610227]. Seborrhea-like dermatitis with psoriasiform is characterized by a chronic fine diffuse scaly erythematous rash on the face, particularly on the chin, nasolabial folds and eyebrows, around earlobes and over the scalp. The rash exacerbate in the winter, with emotional stress and after strenuous physical activity. Hyperkeratosis of skin over the elbows, knees, palms, soles and metacarpophalangeal joints is evident. There is no arthralgia, arthritis or neurological disorders.,tissue specificity:Expressed in the skin, prostate, lungs placenta and thymus, and at low level in T-cells. Not expressed in peripheral blood leukocytes, pancreas and brain. Clearly expressed in primary keratinocytes but not in fibroblasts.,
Background	zinc finger protein 750(ZNF750) Homo sapiens This gene encodes a protein with a nuclear localization site and a C2H2 zinc finger domain. Mutations in this



UpingBio technology Co.,Ltd





gene have been associated with seborrhea-like dermatitis with psoriasiform elements. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images