

(Tel: 400-999-8863 ■ Email:Upingbio.163.com



TTC19 Polyclonal Antibody

5 TPR repeats., Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a		
Reactivity Human;Mouse Applications WB;ELISA Gene Name TTC19 Protein Name Tetratricopeptide repeat protein 19, mitochondrial (TPR repeat protein 19) Immunogen Synthesized peptide derived from part region of human protein Specificity TTC19 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator, similarity Contains 5 TPR repeats. Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPPs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of organisms including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III, It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III (provided by RefSeq, Sep 2012). matters needing Avoid repeated freezing and thawing!	Catalog No	YP-Ab-06511
Applications WB;ELISA Gene Name TTC19 Protein Name Tetratricopeptide repeat protein 19, mitochondrial (TPR repeat protein 19) Immunogen Synthesized peptide derived from part region of human protein Specificity TTC19 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of organisms including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012].	Isotype	IgG
Gene Name TTC19 Protein Name Tetratricopeptide repeat protein 19, mitochondrial (TPR repeat protein 19) Immunogen Synthesized peptide derived from part region of human protein Specificity TTC19 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Brain, Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator., similarity:Contains 5 TPR repeats. Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungl and plants, and aerivolved in a variety of organisms including portein, fungl and plants, and envolved in a variety of organisms including portein, fungl and plants, and envolved in a variety of organisms including bacteria, fung	Reactivity	Human;Mouse
Protein Name Tetratricopeptide repeat protein 19, mitochondrial (TPR repeat protein 19) Immunogen Synthesized peptide derived from part region of human protein Specificity TTC19 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012].	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein Specificity TTC19 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungia and plants, and are involved in a variety of organisms including bacteria, fungia and plants, and are involved in the formation of the mitochondrial respiratory chain III, It has also been suggested that this protein is embeddeded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III, It has also been suggested that this pro	Gene Name	TTC19
Specificity TTC19 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq. Sep 2012], matters needing Avoid repeated freezing and thawing!	Protein Name	Tetratricopeptide repeat protein 19, mitochondrial (TPR repeat protein 19)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain, Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator., similarity: Contains 5 TPR repeats., Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of functions including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interaction. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from part region of human protein
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of or organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Specificity	TTC19 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., Background These or a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Purification	, , , , , , , , , , , , , , , , , , , ,
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane. Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band 41kD Cell Pathway Mitochondrion inner membrane . Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band 41kD Cell Pathway Mitochondrion inner membrane. Tissue Specificity Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Mitochondrion inner membrane . Brain,Muscle, Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Brain,Muscle, caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Observed Band	41kD
Function caution:It is uncertain whether Met-1 or Met-122 is the initiator.,similarity:Contains 5 TPR repeats., This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Cell Pathway	Mitochondrion inner membrane .
Background This gene encodes a protein with a tetratricopeptide repeat (TPR) domain containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], Matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Brain,Muscle,
containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2012], matters needing Avoid repeated freezing and thawing!	Function	
	Background	containing several TPRs of about 34 aa each. These repeats are found in a variety of organisms including bacteria, fungi and plants, and are involved in a variety of functions including protein-protein interactions. This protein is embedded in the inner mitochondrial membrane and is involved in the formation of the mitochondrial respiratory chain III. It has also been suggested that this protein plays a role in cytokinesis. Mutations in this gene cause mitochondrial complex III deficiency. Alternatively spliced transcript variants have been found for this gene.
		Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd






Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night