

**(** Tel: 400-999-8863 ■ Email:Upingbio.163.com





## NUFP2 Polyclonal Antibody

Catalog No	YP-Ab-06517
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	NUFIP2 KIAA1321 PIG1
Protein Name	Nuclear fragile X mental retardation-interacting protein 2 (82 kDa FMRP-interacting protein) (82-FIP) (Cell proliferation-inducing gene 1 protein) (FMRP-interacting protein 2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 180-260
Specificity	NUFP2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	76kD
Observed Band Cell Pathway	76kD  Nucleus . Cytoplasm . Cytoplasm, Stress granule . Distribution is cell cycle-modulated, being cytoplasmic in the G2/M phase and accumulating in nucleus during the G1 phase (PubMed:12837692)
	Nucleus . Cytoplasm . Cytoplasm, Stress granule . Distribution is cell cycle-modulated, being cytoplasmic in the G2/M phase and accumulating in
Cell Pathway	Nucleus . Cytoplasm . Cytoplasm, Stress granule . Distribution is cell cycle-modulated, being cytoplasmic in the G2/M phase and accumulating in nucleus during the G1 phase (PubMed:12837692)
Cell Pathway  Tissue Specificity	Nucleus . Cytoplasm . Cytoplasm, Stress granule . Distribution is cell cycle-modulated, being cytoplasmic in the G2/M phase and accumulating in nucleus during the G1 phase (PubMed:12837692)  Brain, Cervix carcinoma, Epithelium, PCR rescued clones, T-cell, Uterus,  function: Binds RNA., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., subcellular location: Distribution is cell cycle-modulated, being cytoplasmic in the G2/M phase and accumulating in nucleus during the G1
Cell Pathway  Tissue Specificity  Function	Nucleus . Cytoplasm . Cytoplasm, Stress granule . Distribution is cell cycle-modulated, being cytoplasmic in the G2/M phase and accumulating in nucleus during the G1 phase (PubMed:12837692)  Brain, Cervix carcinoma, Epithelium, PCR rescued clones, T-cell, Uterus,  function: Binds RNA., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., subcellular location: Distribution is cell cycle-modulated, being cytoplasmic in the G2/M phase and accumulating in nucleus during the G1 phase., subunit: Interacts with FMRP.,  function: Binds RNA., PTM: Phosphorylated upon DNA damage, probably by ATM or ATR., subcellular location: Distribution is cell cycle-modulated, being cytoplasmic in the G2/M phase and accumulating in nucleus during the G1



## UpingBio technology Co.,Ltd






**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of Hela lysis, using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000