

Tel: 400-999-8863
 ■ Email:Upingbio.163.com





AEBP1 Polyclonal Antibody

Catalog No	YP-Ab-06523
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	AEBP1 ACLP
Protein Name	Adipocyte enhancer-binding protein 1 (AE-binding protein 1) (Aortic carboxypeptidase-like protein)
Immunogen	Synthesized peptide derived from part region of human protein. AA range: 780-810
Specificity	AEBP1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	127kD
Cell Pathway	[Isoform 1]: Secreted .; [Isoform 2]: Cytoplasm . Nucleus .
Tissue Specificity	Expressed in osteoblast and visceral fat.
Function	caution:This protein has lost active site residues and zinc-binding sites and so is unlikely to be catalytically active.,function:May positively regulate MAP-kinase activity in adipocytes, leading to enhanced adipocyte proliferation and reduced adipocyte differentiation (By similarity). May also positively regulate NF-kappa-B activity in macrophages by promoting the phosphorylation and subsequent degradation of I-kappa-B-alpha (NFKBIA), leading to enhanced macrophage
	inflammatory responsiveness (By similarity). Can act as a transcriptional repressor.,PTM:Phosphorylated by MAPK1 in vitro.,similarity:Belongs to the peptidase M14 family.,similarity:Contains 1 F5/8 type C domain.,subunit:Interacts with GNG5, NFKBIA, MAPK1, MAPK3 and PTEN (By similarity). May interact with calmodulin (By similarity). Binds to DNA in vitro.,tissue specificity:Expressed in osteoblast and visceral fat.,
Background	repressor.,PTM:Phosphorylated by MAPK1 in vitro.,similarity:Belongs to the peptidase M14 family.,similarity:Contains 1 F5/8 type C domain.,subunit:Interacts with GNG5, NFKBIA, MAPK1, MAPK3 and PTEN (By similarity). May interact with calmodulin (By similarity). Binds to DNA in vitro.,tissue specificity:Expressed in



UpingBio technology Co.,Ltd

Tel: 400-999-8863
 ■ Email:Upingbio.163.com





adipogenesis and smooth muscle cell differentiation. Studies in mice suggest that this gene functions in wound healing and abdominal wall development.

Overexpression of this gene is associated with glioblastoma. [provided by RefSeq, May 20'13],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images