



XRN1 Polyclonal Antibody

Catalog No	YP-Ab-06531
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	XRN1 SEP1
Protein Name	5'-3' exoribonuclease 1 (EC 3.1.11.-) (Strand-exchange protein 1 homolog)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	XRN1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	187kD
Cell Pathway	Cytoplasm . Discrete foci at the inner surface of the cell membrane.
Tissue Specificity	Expressed in heart, brain, pancreas, spleen, testis, osteogenic sarcoma (OGS) biopsy and primary cell lines.
Function	function:Major 5'-3' exoribonuclease involved in mRNA decay. Required for the 5'-3'processing of the G4 tetraplex-containing DNA and RNA substrates. The kinetic of hydrolysis is faster for G4 RNA tetraplex than for G4 DNA tetraplex and monomeric RNA tetraplex. Binds to RNA and DNA (By similarity). May act as a tumor suppressor protein in osteogenic sarcoma (OGS).,induction:By GDNF.,miscellaneous:Down-regulated in OGS biopsy.,similarity:Belongs to the 5'-3' exonuclease family.,subcellular location:Discrete foci at the inner surface of the cell membrane.,subunit:Associates with alpha and beta tubulins (By similarity). Found in a mRNP complex with RENT1, RENT2, RENT3B and XRN1.,tissue specificity:Expressed in heart, brain, pancreas, spleen, testis, osteogenic sarcoma (OGS) biopsy and primary cell lines.,
Background	This gene encodes a member of the 5'-3' exonuclease family. The encoded protein may be involved in replication-dependent histone mRNA degradation, and interacts directly with the enhancer of mRNA-decapping protein 4. In addition to mRNA metabolism, a similar protein in yeast has been implicated

in a variety of nuclear and cytoplasmic functions, including homologous recombination, meiosis, telomere maintenance, and microtubule assembly. Mutations in this gene are associated with osteosarcoma, suggesting that the encoded protein may also play a role in bone formation. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images