

Tel: 400-999-8863 
■ Email:Upingbio.163.com





## MINA Polyclonal Antibody

| Catalog No         | YP-Ab-06546  |  |
|--------------------|--|--|
| Isotype            | IgG  |  |
| Reactivity         | Human;Rat;Mouse;   |  |
| Applications       | WB;ELISA   |  |
| Gene Name          | MINA MDIG MINA53 NO52  |  |
| Protein Name       | MYC-induced nuclear antigen (Mineral dust-induced gene protein) (Nucleolar protein 52)   |  |
| Immunogen          | Synthesized peptide derived from part region of human protein  |  |
| Specificity        | MINA Polyclonal Antibody detects endogenous levels of protein.   |  |
| Formulation        | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.   |  |
| Source             | Polyclonal, Rabbit,IgG   |  |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |  |
| Dilution           | WB 1:500-2000 ELISA 1:5000-20000   |  |
| Concentration      | 1 mg/ml  |  |
| Purity             | ≥90%   |  |
| Storage Stability  | -20°C/1 year   |  |
| Synonyms           |  |  |
| Observed Band      | 51kD   |  |
| Cell Pathway       | Nucleus . Nucleus, nucleolus .   |  |
| Tissue Specificity | Expressed in liver, skeletal muscle, heart, pancreas, and placenta. Not detected in brain, lung or kidney. Expressed in several lung cancer tissues, but is barely detected in the adjacent non-cancerous tissues. Also highly expressed in several esophageal squamous cell carcinoma (ESCC), and colon cancer tissues, and in various cancer cell lines.   |  |
| Function           | function:Involved in cellular proliferation. May play an important role in cell growth and survival. May be involved in ribosome biogenesis, most likely during the assembly process of pre-ribosomal particles.,induction:Up-regulated in response to MYC, in alveolar macrophages from coal miners and in silica particle-treated A549 lung cancer cells.,sequence caution:Translated as Trp.,similarity:Belongs to the MINA53/NO66 family.,similarity:Contains 1 JmjC domain.,tissue specificity:Expressed in liver, skeletal muscle, heart, pancreas, and placenta. Not detected in brain, lung or kidney. Expressed in several lung cancer tissues, but is barely detected in the adjacent non-cancerous tissues. Also highly expressed in several esophageal squamous cell carcinoma (ESCC), and colon cancer tissues, |  |

and in various cancer cell lines.,



## UpingBio technology Co.,Ltd

🕻 Tel: 400-999-8863 🗷 Emall:Upingbio.163.com



| Background                | MINA is a c-Myc (MYC; MIM 190080) target gene that may play a role in cell proliferation or regulation of cell growth. (Tsuneoka et al., 2002 [PubMed 12091391]; Zhang et al., 2005 [PubMed 15897898]).[supplied by OMIM, May 2008], |  |
|---------------------------|--|--|
| matters needing attention | Avoid repeated freezing and thawing!   |  |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |  |

| Products Images |
|-----------------|
|                 |
|                 |
|                 |
|                 |
|                 |
|                 |